

# Abin-2 (K-17): sc-32638

## BACKGROUND

The nuclear factor NFκB is essential for the regulation of immune response genes, inflammatory processes and apoptosis. Abin-2 (also designated A20-binding inhibitor of NFκB activation 2) is an intracellular zinc-finger protein that inhibits the expression of NFκB by recruiting a chromatin-remodeling complex to the target gene. Abin-2, a p105-associated protein, is a potent inhibitor of TNF-induced cell death. Abin-2 can also associate with TPL-2, and in endogenous tissues it is frequently associated with both TPL-2 and p105. siRNA depletion of Abin-2 has been found to reduce levels of TPL-2 but not of p105, which indicates that Abin-2 is involved in the TLR4 signaling pathway. Abin-2 inhibits endothelial apoptosis, but upon deletion of the carboxy-terminus of the protein, its ability to inhibit apoptosis is removed.

## CHROMOSOMAL LOCATION

Genetic locus: TNIP2 (human) mapping to 4p16.3; Tnip2 (mouse) mapping to 5 B2.

## SOURCE

Abin-2 (K-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Abin-2 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-32638 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-32638 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

Abin-2 (K-17) is recommended for detection of Abin-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Abin-2 (K-17) is also recommended for detection of Abin-2 in additional species, including equine, canine and bovine.

Suitable for use as control antibody for Abin-2 siRNA (h): sc-44638, Abin-2 siRNA (m): sc-44639, Abin-2 shRNA Plasmid (h): sc-44638-SH, Abin-2 shRNA Plasmid (m): sc-44639-SH, Abin-2 shRNA (h) Lentiviral Particles: sc-44638-V and Abin-2 shRNA (m) Lentiviral Particles: sc-44639-V.

Abin-2 (K-17) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of Abin-2 isoforms: 49/37 kDa.

Positive Controls: SW480 cell lysate: sc-2219.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



Abin-2 (K-17): sc-32638. Immunoperoxidase staining of formalin fixed, paraffin-embedded human thyroid gland tissue showing nuclear staining of glandular cells.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

**MONOS**  
Satisfaction  
Guaranteed

Try **Abin-2 (H-8): sc-271850**, our highly recommended monoclonal alternative to Abin-2 (K-17).