

DEC2 (H-72): sc-32853

BACKGROUND

Human DEC1 is a 412 amino acid, basic helix-loop-helix (bHLH) containing protein that is involved in the control of proliferation and/or differentiation of several cell types including nerve cells, fibroblasts and chondrocytes. The bHLH region of DEC1 is structurally similar to the bHLH regions of the mammalian HES family, *Drosophila* hairy and enhancer of split m7. DEC1 is a novel direct target for cAMP in a wide range of cells and is involved in the control of gene expression in cAMP-activated cells. DEC2, also known as SHARP1, is highly expressed in skeletal muscle and brain. The gene encoding human DEC2 maps to chromosome 12p12.1. DEC1 and DEC2 play a role in regulating the mammalian molecular clock by suppressing the transcription of specific clock genes. Both DEC1 and DEC2 are detected in the suprachiasmatic nucleus in a circadian fashion. Brief light impulses induce the expression of DEC1 in a phase-dependent manner.

REFERENCES

1. Shen, M., et al. 1997. Molecular characterization of the novel basic helix-loop-helix protein DEC1 expressed in differentiated human embryo chondrocytes. *Biochem. Biophys. Res. Commun.* 236: 294-298.
2. Shen, M., et al. 2001. Induction of basic helix-loop-helix protein DEC1 (bHLHB2)/Stra13/SHARP2 in response to the cyclic adenosine monophosphate pathway. *Eur. J. Cell Biol.* 80: 329-334.
3. Fujimoto, K., et al. 2001. Molecular cloning and characterization of DEC2, a new member of basic helix-loop-helix proteins. *Biochem. Biophys. Res. Commun.* 280: 164-171.
4. Honma, S., et al. 2002. DEC1 and DEC2 are regulators of the mammalian molecular clock. *Nature* 419: 841-844.
5. LocusLink Report (LocusID: 8553). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: BHLHE41 (human) mapping to 12p12.1; Bhlhe41 (mouse) mapping to 6 G3.

SOURCE

DEC2 (H-72) is a rabbit polyclonal antibody raised against amino acids 411-482 mapping at the C-terminus of DEC2 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-32853 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

DEC2 (H-72) is recommended for detection of DEC2 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

DEC2 (H-72) is also recommended for detection of DEC2 in additional species, including porcine.

Suitable for use as control antibody for DEC2 siRNA (h): sc-37769, DEC2 siRNA (m): sc-37770, DEC2 shRNA Plasmid (h): sc-37769-SH, DEC2 shRNA Plasmid (m): sc-37770-SH, DEC2 shRNA (h) Lentiviral Particles: sc-37769-V and DEC2 shRNA (m) Lentiviral Particles: sc-37770-V.

DEC2 (H-72) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

1. Imaizumi, T., et al. 2011. Basic-helix-loop-helix transcription factor DEC2 constitutes negative feedback loop in IFN-β-mediated inflammatory responses in human mesangial cells. *Immunol. Lett.* 136: 37-43.

RESEARCH USE

For research use only, not for use in diagnostic procedures.


 MONOS
 Satisfation
 Guaranteed

Try **DEC2 (E-4): sc-373763**, our highly recommended monoclonal alternative to DEC2 (H-72).