

Ebi3 (H-110): sc-32868

BACKGROUND

Epstein-Barr virus-induced gene 3 (Ebi3) is a widely expressed homolog to the interleukin IL-12p40 subunit protein that forms a heterodimer with either IL-12p35 or an IL-12p35 homologue, p28, to create a new cytokine (IL-27). IL-27 is an early product of activated antigen-presenting cells and drives rapid clonal expansion of naive but not memory CD4⁺ T cells. Interferon- β differentially regulates expression of the IL-12 family members p35, p40, p19 and Ebi3 in activated human dendritic cells. Ebi3 may function to antagonize IL-12 and to inhibit the development of a Th1 immune response. Ebi3 is strongly expressed in Hodgkin and Reed-Sternberg cells, independently of the EBV status of the tumor cells. Research suggest that Ebi3 may be an additional component of the repertoire employed by Hodgkin and Reed-Sternberg cells to inhibit an effective anti-tumor or anti-viral immune response. The human Ebi3 gene maps to chromosome 19p13.3 and encodes a secreted glycoprotein that is expressed in the spleen and tonsils, and at high levels in full-term placenta.

REFERENCES

1. Devergne, O., et al. 2001. Expression of Epstein-Barr virus-induced gene 3, an interleukin-12 p40-related molecule, throughout human pregnancy: involvement of syncytiotrophoblasts and extravillous trophoblasts. *Am. J. Pathol.* 159: 1763-1776.
2. Omata, F., et al. 2001. The expression of IL-12 p40 and its homologue, Epstein-Barr virus-induced gene 3, in inflammatory bowel disease. *Inflamm. Bowel Dis.* 7: 215-220.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 605816. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Nieuwenhuis, E.E., et al. 2002. Disruption of T helper 2-immune responses in Epstein-Barr virus-induced gene 3-deficient mice. *Proc. Natl. Acad. Sci. USA* 99: 16951-16956.
5. Niedobitek, G., et al. 2002. Frequent expression of the Epstein-Barr virus (EBV)-induced gene, EBI3, an IL-12 p40-related cytokine, in Hodgkin and Reed-Sternberg cells. *J. Pathol.* 198: 310-316.

CHROMOSOMAL LOCATION

Genetic locus: EBI3 (human) mapping to 19p13.3; Ebi3 (mouse) mapping to 17 C.

SOURCE

Ebi3 (H-110) is a rabbit polyclonal antibody raised against amino acids 21-130 mapping near the N-terminus of Ebi3 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Ebi3 (H-110) is recommended for detection of Ebi3 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for Ebi3 siRNA (h): sc-39410, Ebi3 siRNA (m): sc-39411, Ebi3 shRNA Plasmid (h): sc-39410-SH, Ebi3 shRNA Plasmid (m): sc-39411-SH, Ebi3 shRNA (h) Lentiviral Particles: sc-39410-V and Ebi3 shRNA (m) Lentiviral Particles: sc-39411-V.

Molecular Weight of Ebi3: 34 kDa.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



Ebi3 (H-110): sc-32868. Western blot analysis of Ebi3 expression in I-11.15 whole cell lysate.



Ebi3 (H-110): sc-32868. Western blot analysis of Ebi3 expression in RAW 264.7 whole cell lysate.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

MONOS
Satisfaction
Guaranteed

Try **Ebi3 (G-4): sc-166158** or **Ebi3 (F-7): sc-515323**, our highly recommended monoclonal alternatives to Ebi3 (H-110).