

ADRP (H-80): sc-32888

BACKGROUND

Mannose 6-phosphate receptors (MPRs) deliver lysosomal hydrolase to endosomes from the Golgi and back again. Cargo selection protein TIP47, also designated placental protein 17, is required for the transport from endosomes to the *trans*-Golgi network and interacts with the cytoplasmic domains of both cation-dependent and cation-independent MPRs. Another member of the perilipin family, Adipophilin (ADRP), is a protein associated with the globule surface membrane material of milk lipid globules. The phosphoprotein Perilipin (Peri) is located on the surface of intracellular lipid droplets within adipocytes where it protects lipid storage droplets by coating them in adipocytes until they are digested by lipase. As a critical regulator of lipolysis, elevated Perilipin levels have been linked to obesity.

REFERENCES

1. Heid, H.W., et al. 1996. Adipocyte differentiation-related protein is secreted into milk as a constituent of milk lipid globule membrane. *Biochem. J.* 320: 1025-1030.
2. Souza, S.C., et al. 2002. Modulation of hormone-sensitive lipase and protein kinase A-mediated lipolysis by perilipin A in an adenoviral reconstituted system. *J. Biol. Chem.* 277: 8267-8272.
3. Kern, P.A., et al. 2004. Perilipin expression in human adipose tissue is elevated with obesity. *J. Clin. Endocrinol. Metab.* 89: 1352-1358.
4. Elchalal, U., et al. 2005. Insulin and fatty acids regulate the expression of the fat droplet-associated protein adipophilin in primary human trophoblasts. *Am. J. Obstet. Gynecol.* 193: 1716-1723.

CHROMOSOMAL LOCATION

Genetic locus: ADFP (human) mapping to 9p22.1; Adfp (mouse) mapping to 4 C4.

SOURCE

ADRP (H-80) is a rabbit polyclonal antibody raised against amino acids 321-400 mapping near the C-terminus of ADRP of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

ADRP (H-80) is recommended for detection of ADRP of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

ADRP (H-80) is also recommended for detection of ADRP in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for ADRP siRNA (h): sc-44841, ADRP siRNA (m): sc-44842, ADRP shRNA Plasmid (h): sc-44841-SH, ADRP shRNA Plasmid (m): sc-44842-SH, ADRP shRNA (h) Lentiviral Particles: sc-44841-V and ADRP shRNA (m) Lentiviral Particles: sc-44842-V.

Molecular Weight of ADRP: 48 kDa.

Positive Controls: human liver extract: sc-363766.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

SELECT PRODUCT CITATIONS

1. Sakurai, R., et al. 2011. Curcumin augments lung maturation, preventing neonatal lung injury by inhibiting TGF-β signaling. *Am. J. Physiol. Lung Cell Mol. Physiol.* 301: L721-L730.
2. Gandolfi, G., et al. 2011. Perilipin 1 and perilipin 2 protein localization and gene expression study in skeletal muscles of European cross-breed pigs with different intramuscular fat contents. *Meat Sci.* 88: 631-637.
3. Coín Aragüez, L., et al. 2013. Thymus fat as an attractive source of angiogenic factors in elderly subjects with myocardial ischemia. *Age* 35: 1263-1275.

MONOS
Satisfaction
Guaranteed

Try **ADRP (B-6): sc-377429**, our highly recommended monoclonal alternative to ADRP (H-80). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **ADRP (B-6): sc-377429**.