

p-PDPK1 (Ser 241): sc-32960

BACKGROUND

PDPK1 (3-phosphoinositide dependent protein kinase 1), also known as PDK1, PDPK2, PDPK2P or PRO0461, is 556 amino acid ubiquitously expressed protein that localizes to the cell membrane, cytoplasm and nucleus. Acting as a master kinase, PDPK1 phosphorylates and activates a subgroup of the AGC family of protein kinases. PDPK1 is involved in mediating signal transduction for controlling proliferation, survival, and growth of developing pancreatic beta cells, regulating Ca²⁺ uptake and Ca²⁺-activated K⁺ channels of mast cells, regulation of chemotaxis and motility of vascular endothelial cells, cardiac homeostasis, and thymocyte development. Belonging to the protein kinase superfamily, PDPK1 contains a PH domain, which play an essential role in homodimerization, localization and nuclear import of PDPK1, and a protein kinase domain. PDPK1 exists as five alternatively spliced isoforms and is encoded by a gene located on human chromosome 16p13.3.

REFERENCES

- Burgering, B.M., et al. 1995. Protein kinase B (c-Akt) in phosphatidylinositol-3-OH kinase signal transduction. *Nature* 376: 599-602.
- Cross, D.A., et al. 1995. Inhibition of glycogen synthase kinase-3 by Insulin mediated by protein kinase B. *Nature* 378: 785-789.
- Wada, H., et al. 1995. Rejection antigen peptides on BALB/CRL male 1 leukemia recognized by cytotoxic T lymphocytes: derivation from the normally untranslated 5' region of the c-Akt proto-oncogene activated by long terminal repeat. *Cancer Res.* 55: 4780-4783.
- Datta, K., et al. 1995. AH/PH domain-mediated interaction between Akt molecules and its potential role in Akt regulation. *Mol. Cell. Biol.* 15: 2304-2310.
- Franke, T.F., et al. 1995. The protein kinase encoded by the Akt proto-oncogene is a target of the PDGF-activated phosphatidylinositol 3-kinase. *Cell* 81: 727-736.
- Cheng, J.Q., et al. 1996. Amplification of Akt2 in human pancreatic cancer cells and inhibition of Akt2 expression and tumorigenicity by antisense RNA. *Proc. Natl. Acad. Sci. USA* 93: 3636-3641.
- Stephens, L., et al. 1998. Protein kinase B kinases that mediate phosphatidylinositol 3,4,5-trisphosphate-dependent activation of protein kinase B. *Science* 279: 710-714.
- Casamayor, A., et al. 1999. Phosphorylation of Ser 241 is essential for the activity of 3-phosphoinositide-dependent protein kinase-1: identification of five sites of phosphorylation *in vivo*. *Biochem. J.* 342287-342292.

SOURCE

p-PDPK1 (Ser 241) is a rabbit polyclonal antibody raised against a short amino acid sequence containing Ser 241 phosphorylated of PDPK1 of human origin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-32960 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

p-PDPK1 (Ser 241) is recommended for detection of Ser 241 phosphorylated PDPK1 and correspondingly phosphorylated MEK-1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


p-PDPK1 (Ser 241) is also recommended for detection of Ser 241 phosphorylated PDPK1 and correspondingly phosphorylated MEK-1 and 2 in additional species, including equine, canine, bovine and avian.

Molecular Weight of p-PDPK1: 60 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto B Blocking Reagent: sc-2335 (use 50 mM NaF, sc-24988, as diluent), Western Blotting Luminol Reagent: sc-2048 and Lambda Phosphatase: sc-200312A. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

DATA



p-PDPK1 (Ser 241): sc-32960. Immunoperoxidase staining of formalin fixed, paraffin-embedded human placenta tissue showing membrane, cytoplasmic and nuclear staining of decidual cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.