

HoxD10 (E-20): sc-33005

BACKGROUND

The Hox proteins play a role in development and cellular differentiation by regulating downstream target genes. Specifically, the Hox proteins direct DNA-protein and protein-protein interactions that assist in determining the morphologic features associated with the anterior-posterior body axis. Hox proteins are involved in controlling axial patterning, leukemias and hereditary malformations. Homeobox protein HoxD10, also designated Hox-4D or Hox-4E, belongs to the Abd-B homeobox family of proteins. HoxD10 is a nuclear protein primarily expressed in the adult male and female urogenital tracts but also expressed in developing limb buds during development. Defects in the gene encoding for the HoxD10 protein cause congenital vertical talus (CVT), more commonly known as rocker-bottom foot deformity. CVT is characterized by a dislocation of the talonavicular joint but is usually accompanied by other congenital deformities.

REFERENCES

1. Zappavigna, V., et al. 1991. HOX4 genes encode transcription factors with potential auto- and cross-regulatory capacities. *EMBO J.* 10: 4177-4187.
2. Redline, R.W., et al. 1992. Human HOX4E: a gene strongly expressed in the adult male and female urogenital tracts. *Genomics* 13: 425-430.
3. Gabellini, D., et al. 2003. Early mitotic degradation of the homeoprotein HoxC10 is potentially linked to cell cycle progression. *EMBO J.* 22: 3715-3724.
4. Juan, A.H., et al. 2003. Enhancer timing of Hox gene expression: deletion of the endogenous HoxC8 early enhancer. *Development* 130: 4823-4834.
5. Miller, G.J., et al. 2003. Aberrant HoxC expression accompanies the malignant phenotype in human prostate. *Cancer Res.* 63: 5879-5888.
6. Nicolas, S., et al. 2003. The spatial restrictions of 5' HoxC genes expression are maintained in adult newt spinal cord. *Biol. Cell* 95: 589-594.
7. Akbas, G.E. et al. 2004. HOXC and HOXD gene expression in human endometrium: lack of redundancy with HOXA paralogs. *Biol. Reprod.* 70: 39-45.

CHROMOSOMAL LOCATION

Genetic locus: HOXD10 (human) mapping to 2q31.1; Hoxd10 (mouse) mapping to 2 C3.

SOURCE

HoxD10 (E-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of HoxD10 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-33005 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-33005 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

HoxD10 (E-20) is recommended for detection of HoxD10 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

HoxD10 (E-20) is also recommended for detection of HoxD10 in additional species, including equine, canine, bovine, porcine and avian.


Suitable for use as control antibody for HoxD10 siRNA (h): sc-44814, HoxD10 siRNA (m): sc-44815, HoxD10 shRNA Plasmid (h): sc-44814-SH, HoxD10 shRNA Plasmid (m): sc-44815-SH, HoxD10 shRNA (h) Lentiviral Particles: sc-44814-V and HoxD10 shRNA (m) Lentiviral Particles: sc-44815-V.

HoxD10 (E-20) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of HoxD10: 40 kDa.

Positive Controls: HoxD10 (m2): 293T Lysate: sc-126972.

DATA



HoxD10 (E-20): sc-33005. Western blot analysis of HoxD10 expression in non-transfected: sc-117752 (A) and mouse HoxD10 transfected: sc-126972 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

1. Chen, A., et al. 2009. Endothelial cell migration and vascular endothelial growth factor expression are the result of loss of breast tissue polarity. *Cancer Res.* 69: 6721-6729.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **HoxD10 (G-3): sc-166235** or **HoxD10 (B-10): sc-166233**, our highly recommended monoclonal alternatives to HoxD10 (E-20).