

# FBP1 (Q-13): sc-33030

## BACKGROUND

Activation of FUSE, the far-upstream element, is required for the proper expression of the mammalian gene c-Myc in undifferentiated cells. The binding of FBP (FUSE-binding protein or Far upstream element binding protein) to FUSE is necessary for c-Myc expression, indicating that FBP functions as a growth-dependent regulator of c-Myc expression. Isolated from proliferating HL60 cells, FBP, FBP2 and FBP3 comprise a family of single-stranded DNA-binding proteins that specifically bind to FUSE elements. The FBP transcription factors share a conserved central DNA-binding domain and show significant homology in their carboxyl-terminal activation domains. Expression of FBP is detected in undifferentiated cells and is substantially decreased following cellular differentiation.

## CHROMOSOMAL LOCATION

Genetic locus: FUBP1 (human) mapping to 1p31.1; Fubp1 (mouse) mapping to 3 H3.

## SOURCE

FBP1 (Q-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of FBP1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33030 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-33030 X, 200 µg/0.1 ml.

## APPLICATIONS

FBP1 (Q-13) is recommended for detection of FBP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for FBP1 siRNA (h): sc-43760, FBP1 siRNA (m): sc-44829, FBP1 shRNA Plasmid (h): sc-43760-SH, FBP1 shRNA Plasmid (m): sc-44829-SH, FBP1 shRNA (h) Lentiviral Particles: sc-43760-V and FBP1 shRNA (m) Lentiviral Particles: sc-44829-V.

FBP1 (Q-13) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.

Molecular Weight of FBP1: 74 kDa.

Positive Controls: FBP1 (h2): 293T Lysate: sc-171763, HeLa whole cell lysate: sc-2200 or HL-60 whole cell lysate: sc-2209.


## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



FBP1 (Q-13): sc-33030. Western blot analysis of FBP1 expression in non-transfected: sc-117752 (A) and human FBP1 transfected: sc-171763 (B) 293T whole cell lysates.

## SELECT PRODUCT CITATIONS

- Jang, M., et al. 2009. Far upstream element-binding protein-1, a novel caspase substrate, acts as a cross-talker between apoptosis and the c-myc oncogene. *Oncogene* 28: 1529-1536.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **FBP1 (G-8): sc-271241** or **FBP1 (A-4): sc-393928**, our highly recommended monoclonal alternatives to FBP1 (Q-13).