

RB3 (C-12): sc-33071

BACKGROUND

Op18 (for oncoprotein 18, also designated Stathmin, prosolin or metablastin) is a conserved, tubulin-associated, intracellular protein. It serves as a transducing protein, via phosphorylation, for a variety of cell signaling pathways and is involved in both mitosis and differentiation. Op18 is present in many cancers, including breast carcinoma and different leukemias. The neuronal growth-associated protein (nGAP) Stathmin-2, which shares sequence homology with the phosphoprotein Op18/Stathmin-1, is expressed in a variety of neural, immune and reproductive system cell types. Stathmin-2 gene expression is altered in age-related neurodegenerative diseases such as Alzheimer's disease. Stathmin-4 (also designated Stathmin-like protein B3 or RB3) is a Stathmin-like protein involved in the destabilization of microtubules, specifically in brain tissue. RB3 has a unique N-terminal membrane-associated domain and a Stathmin-like domain at the C-terminus. This C-terminal domain promotes microtubule destabilization and tubulin sequestering.

REFERENCES

1. Nakao, C., Itoh, T.J., Hotani, H. and Mori, N. 2004. Modulation of the Stathmin-like microtubule destabilizing activity of RB3, a neuron-specific member of the SCG10 family, by its N-terminal domain. *J. Biol. Chem.* 279: 23014-23021.
2. Iancu-Rubin, C., Nasrallah, C.A. and Atweh, G.F. 2005. Stathmin prevents the transition from a normal to an endomitotic cell cycle during megakaryocytic differentiation. *Cell Cycle* 4: 1774-1782.
3. Shumyatsky, G.P., Malleret, G., Shin, R.M., Takizawa, S., Tully, K., Tsvetkov, E., Zakharenko, S.S., Joseph, J., Vronskaya, S., Yin, D., Schubart, U.K., Kandel, E.R. and Bolshakov, V.Y. 2005. Stathmin, a gene enriched in the amygdala, controls both learned and innate fear. *Cell* 123: 697-709.
4. Giampietro, C., Luzzati, F., Gambarotta, G., Giacobini, P., Boda, E., Fasolo, A. and Perroteau, I. 2005. Stathmin expression modulates migratory properties of GN-11 neurons *in vitro*. *Endocrinology* 146: 1825-1834.

CHROMOSOMAL LOCATION

Genetic locus: STMN4 (human) mapping to 8p21.2; Stmn4 (mouse) mapping to 14 D1.

SOURCE

RB3 (C-12) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of RB3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33071 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

RB3 (C-12) is recommended for detection of RB3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for RB3 siRNA (h): sc-40784, RB3 siRNA (m): sc-40785, RB3 shRNA Plasmid (h): sc-40784-SH, RB3 shRNA Plasmid (m): sc-40785-SH, RB3 shRNA (h) Lentiviral Particles: sc-40784-V and RB3 shRNA (m) Lentiviral Particles: sc-40785-V.


Molecular Weight of RB3: 29 kDa.

Positive Controls: mouse brain extract: sc-2253.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



RB3 (C-12): sc-33071. Western blot analysis of RB3 expression in mouse brain tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

MONOS
Satisfaction
Guaranteed

Try **RB3 (B-11): sc-376829** or **RB3 (B-3): sc-376936**, our highly recommended monoclonal alternatives to RB3 (C-12).