

PP2B-B1/2 (FL-170): sc-33166

BACKGROUND

In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases. In general, the protein phosphatase (PP) holoenzyme is a trimeric complex composed of a regulatory subunit, a variable subunit and a catalytic subunit. Four major families of protein phosphatase catalytic subunit have been identified, designated PP1, PP2A, PP2B and PP2C. An additional protein phosphatase catalytic subunit, PPX (also known as PP4), is a putative member of a novel PP family. The PP2B family comprises subfamily members PP2B-A α , PP2B-A β and PP2B-A γ . Two additional regulatory subunits been identified, designated PP2B-B1 and PP2B-B2.

REFERENCES

1. Ueki, K., et al. 1992. Structure and expression of two isoforms of the murine calmodulin-dependent protein phosphatase regulatory subunit (calcineurin B). *Biochem. Biophys. Res. Commun.* 187: 537-543.
2. Mumby, M.C., et al. 1993. Protein serine/threonine phosphatases: structure, regulation and functions in cell growth. *Physiol. Rev.* 73: 673-699.
3. Hendrix, P., et al. 1993. Structure and expression of a 72 kDa regulatory subunit of protein phosphatase 2A. Evidence for different size forms produced by alternative splicing. *J. Biol. Chem.* 268: 15267-15276.
4. Cohen, P.T. 1993. Important roles for novel protein phosphatases dephosphorylating serine and threonine residues. *Biochem. Soc. Trans.* 21: 884-888.
5. Okubo, S., et al. 1994. A regulatory subunit of smooth muscle myosin bound phosphatase. *Biochem. Biophys. Res. Commun.* 200: 429-434.
6. Wera, S., et al. 1995. Serine/threonine protein phosphatases. *Biochem. J.* 311: 17-29.
7. Lohse, D.L., et al. 1995. Insights derived from the structures of the Ser/Thr phosphatases calcineurin and protein phosphatase 1. *Structure* 3: 987-990.

CHROMOSOMAL LOCATION

Genetic locus: PPP3R1 (human) mapping to 2p14, PPP3R2 (human) mapping to 9q31.1; Ppp3r1 (mouse) mapping to 11 A2, Ppp3r2 (mouse) mapping to 4 B1.

SOURCE

PP2B-B1/2 (FL-170) is a rabbit polyclonal antibody raised against amino acids 1-170 representing full length PP2B-B2 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

PP2B-B1/2 (FL-170) is recommended for detection of PP2B-B1 and PP2B-B2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of PP2B-B1: 19 kDa.


Molecular Weight of PP2B-B2: 21 kDa.

Positive Controls: Human PP2B-B1/2 transfected HEK293T whole cell lysate.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz MarkerTM compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz MarkerTM Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruzTM Mounting Medium: sc-24941.

DATA



PP2B-B1/2 (FL-170): sc-33166. Western blot analysis of PP2B-B1/2 expression in non-transfected (A) and human PP2B-B1/2 transfected (B) HEK293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **PP2B-B1/2 (D-1): sc-373803** or **PP2B-B1 (70-A): sc-130393**, our highly recommended monoclonal alternatives to PP2B-B1/2 (FL-170).