

SRY (M-140): sc-33169

BACKGROUND

SRY (sex-determining region Y protein) is a transcriptional activator required for male sex determination in mammals. This protein, also referred to as testis-determining factor (TDF), is an HMG box protein that initiates the formation of testis from undifferentiated gonad. The DNA-binding activity of SRY is required for normal testis formation. This DNA-binding activity is thought to be regulated by PKA, which phosphorylates SRY *in vivo*. Mutations in SRY have been associated with 46,XY gonadal dysgenesis, in which the gonads fail to develop in XY phenotypic females.

REFERENCES

1. Clepet, C., et al. 1993. The human SRY transcript. *Hum. Mol. Genet.* 2: 2007-2012.
2. Harley, V.R., et al. 1994. The biochemical role of SRY in sex determination. *Mol. Reprod. Dev.* 39: 184-193.
3. Fechner, P.Y. 1996. The role of SRY in mammalian sex determination. *Acta Paediatr. Jpn.* 38: 380-389.
4. Tsutsumi, O., et al. 1996. Analysis of the testis-determining gene SRY in patients with gonadal dysgenesis. *Horm. Res.* 46: 6-10.
5. Graves, J.A. 1998. Evolution of the mammalian Y chromosome and sex-determining genes. *J. Exp. Zool.* 281: 472-481.

SOURCE

SRY (M-140) is a rabbit polyclonal antibody raised against amino acids 1-140 mapping at the N-terminus of SRY of mouse origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-33169 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

SRY (M-140) is recommended for detection of SRY and SOX family proteins of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SRY (M-140) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of SRY: 27 kDa.

Positive Controls: mouse testis extract: sc-2405.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



SRY (M-140): sc-33169. Western blot analysis of SRY expression in mouse testis tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
Satisfaction
Guaranteed

Try **SRY (D-11): sc-398567**, our highly recommended monoclonal alternative to SRY (M-140).