

# BMP-1 (H-300): sc-33200

## BACKGROUND

Bone morphogenic proteins (BMPs) are members of the TGF $\beta$  superfamily. BMPs are involved in the induction of cartilage and bone formation. *In vivo* studies have shown that BMP-2 (also designated BMP-2A) and BMP-3 can independently induce cartilage formation. BMP-1 is a metalloprotease that plays important roles in regulating the deposition of fibrous extracellular matrix in vertebrates, including provision of the procollagen C-proteinase activity that processes the major fibrillar collagens I-III. Smad3 association with the TGF $\beta$  receptor complex and Smad1 translocation to the nucleus are observed after the addition of BMP-4 (also designated BMP-2B), suggesting that BMP-4 may play a role in activation of the Smad pathway. BMP-5, BMP-6 and BMP-7 all share high sequence homology with BMP-2, indicating that they each may be able to induce cartilage formation. BMP-8 (also designated OP-2) is thought to be involved in early development, as detectable expression has not been found in adult organs. Human BMP-1 splice variants occur between residues 703 and 986 and include: BMP1-1, BMP1-4, BMP1-5, BMP1-6, BMP1-7.

## REFERENCES

1. Wozney, J.M., et al. 1988. Novel regulators of bone formation: molecular clones and activities. *Science* 242: 1528-1534.
2. Massague, J. 1990. The transforming growth factor  $\beta$  family. *Ann. Rev. Cell Biol.* 6: 597-641.
3. Oskaynak, E., et al. 1992. Osteogenic protein-2. A new member of the transforming growth factor  $\beta$  superfamily expressed early in embryogenesis. *J. Biol. Chem.* 267: 25220-25227.
4. Gitelman, S.E., et al. 1994. Recombinant Vgr-1/BMP-6-expressing tumors induce fibrosis and endochondral bone formation *in vivo*. *J. Cell Biol.* 126: 1595-1609.

## SOURCE

BMP-1 (H-300) is a rabbit polyclonal antibody raised against amino acids 687-986 mapping at the C-terminus of BMP-1 of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

## APPLICATIONS

BMP-1 (H-300) is recommended for detection of precursor and mature BMP-1 and Tolloid-like protein 1 and 2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

BMP-1 (H-300) is also recommended for detection of precursor and mature BMP-1 and Tolloid-like protein 1 and 2 in additional species, including canine, bovine and porcine.


Molecular Weight of BMP-1 calculated splice variants: 32-98 kDa.

Positive Controls: NIH/3T3 whole cell lysate: sc-2210.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2051 or ABC: sc-2018 rabbit IgG Staining Systems.

## DATA



BMP-1 (H-300): sc-33200. Western blot analysis of BMP-1 expression in NIH/3T3 whole cell lysate.



BMP-1 (H-300): sc-33200. Immunoperoxidase staining of formalin fixed, paraffin-embedded human duodenum tissue showing cytoplasmic staining of glandular cells.

## SELECT PRODUCT CITATIONS

1. Visigalli, D., et al. 2010. Hind limb unloading of mice modulates gene expression at the protein and mRNA level in mesenchymal bone cells. *BMC Musculoskelet. Disord.* 11: 147.