

Keratocan (D-20): sc-33242

BACKGROUND

Small leucine-rich proteoglycans (SLRPs) such as Decorin, Biglycan, Fibromodulin, Keratocan, Lumican and Osteoglycin mediate extracellular matrix organization and are binding partners of TGF β . The Decorin core protein binds to growth factors, intercellular matrix molecules such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Fibromodulin is a collagen-binding keratan sulphate proteoglycan that influences adhesion processes of connective tissue and plays a role in fibrillogenesis by regulating Collagen fibril spacing and thickness. Keratocan develops corneal transparency and maintains the stromal matrix structure. Keratocan (KTN) is a secreted protein in the extracellular matrix that binds to keratan sulfate chains. Keratocan is mainly detected in the cornea, but can also be expressed in trachea, intestine, ovary, lung and skeletal muscle. Defects in the gene encoding for Keratocan can cause cornea plana 2 (CNA2), which is an autosomal recessive disorder where the forward convex curvature of the cornea is flattened.

REFERENCES

1. Tasheva, E.S., et al. 1999. Structure and sequence of the gene encoding human Keratocan. *DNA Seq.* 10: 67-74.
2. Lehmann, O.J., et al. 2001. A novel Keratocan mutation causing autosomal recessive cornea plana. *Invest. Ophthalmol Vis. Sci.* 42: 3118-3122.
3. Wentz-Hunter, K., et al. 2001. Keratocan expression is increased in the stroma of keratoconus corneas. *Mol. Med.* 7: 470-477.
4. Carlson, E.C., et al. 2005. Keratocan, a cornea-specific keratan sulfate proteoglycan, is regulated by Lumican. *J. Biol. Chem.* 280: 25541-25547.
5. Kawakita, T., et al. 2005. Keratocan expression of murine keratocytes is maintained on amniotic membrane by down-regulating TGF β signaling. *J. Biol. Chem.* 280: 27085-27092.
6. Ebenezer, N.D., et al. 2005. Clinical and molecular characterization of a family with autosomal recessive cornea plana. *Arch. Ophthalmol.* 123: 1248-1253.

CHROMOSOMAL LOCATION

Genetic locus: KERA (human) mapping to 12q21.33; Kera (mouse) mapping to 10 C3.

SOURCE

Keratocan (D-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Keratocan of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33242 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4 $^{\circ}$ C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

Keratocan (D-20) is recommended for detection of Keratocan of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Keratocan (D-20) is also recommended for detection of Keratocan in additional species, including equine, canine, bovine and porcine.

Suitable for use as control antibody for Keratocan siRNA (h): sc-44701, Keratocan siRNA (m): sc-44702, Keratocan shRNA Plasmid (h): sc-44701-SH, Keratocan shRNA Plasmid (m): sc-44702-SH, Keratocan shRNA (h) Lentiviral Particles: sc-44701-V and Keratocan shRNA (m) Lentiviral Particles: sc-44702-V.


Molecular Weight of Keratocan: 50 kDa.

Positive Controls: mouse skeletal muscle extract: sc-364250 or JEG-3 whole cell lysate: sc-364255.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



Keratocan (D-20): sc-33242. Western blot analysis of Keratocan expression in mouse skeletal muscle tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.