



## Tap42 (yE-15): sc-33284

### BACKGROUND

In *Saccharomyces cerevisiae*, the target of rapamycin (Tor) pathway, mediates cell proliferation and growth. Two A phosphatase associated protein (Tap42) is part of the Tor signalling pathway and is involved in transcriptional modulation. Tap42 is phosphorylated by Tor and interacts with the catalytic subunits of protein phosphatase 2A (PP2A) and closely related phosphatase Sit4 via their N-terminal domains. Tap42 can also interact with Pph3 and Ppg1, two 2A-like phosphatases. Tap42 acts as an inhibitor of PP2A phosphatase, and the complex between Tap42 and the catalytic subunit of PP2A acts via a Rho GTPase-dependent mechanism to regulate the actin cytoskeleton in *S. cerevisiae*. Upon treatment with rapamycin, Tap42 interacts with TIP41, which binds to and inhibits Tap42. In mammals, the homolog of Tap42 is known as  $\alpha 4$ .

### REFERENCES

- Di Como, C.J., et al. 1996. Nutrients, via the Tor proteins, stimulate the association of Tap42 with type 2A phosphatases. *Genes Dev.* 10: 1904-1916.
- Nanahoshi, M., et al. 1998. Regulation of protein phosphatase 2A catalytic activity by  $\alpha 4$  protein and its yeast homolog Tap42. *Biochem. Biophys. Res. Commun.* 251: 520-526.
- Beck, T., et al. 1999. The Tor signalling pathway controls nuclear localization of nutrient-regulated transcription factors. *Nature* 402: 689-92.
- Jiang, Y., et al. 1999. Tor proteins and protein phosphatase 2A reciprocally regulate Tap42 in controlling cell growth in yeast. *EMBO J.* 18: 2782-2792.
- Cutler, N.S., et al. 2001. The Tor signal transduction cascade controls cellular differentiation in response to nutrients. *Mol. Biol. Cell* 12: 4103-4113.
- Jacinto, E., et al. 2001. TIP41 interacts with Tap42 and negatively regulates the TOR signaling pathway. *Mol. Cell* 8: 1017-1026.
- Torres, J., et al. 2002. Regulation of the cell integrity pathway by rapamycin-sensitive TOR function in budding yeast. *J. Biol. Chem.* 277: 43495-43504.
- Zabrocki, P., et al. 2002. Protein phosphatase 2A on track for nutrient-induced signalling in yeast. *Mol. Microbiol.* 43: 835-842.
- Cherkasova, V.A., et al. 2003. Translational control by Tor and Tap42 through dephosphorylation of eIF2 $\alpha$  kinase GCN2. *Genes Dev.* 17: 859-872.

### SOURCE

Tap42 (yE-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Tap42 of *Saccharomyces cerevisiae* origin.

### PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33284 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### APPLICATIONS

Tap42 (yE-15) is recommended for detection of Tap42 of *Saccharomyces cerevisiae* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Tap42: 42 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.