

SNAT3 (N-20): sc-33445

BACKGROUND

The sodium-coupled neutral amino acid transporters (SNAT) of the SLC38 gene family include system A subtypes SNAT1, SNAT2 and SNAT4 and system N subtypes SNAT3 and SNAT5. The SLC38 transporters are essential for the uptake of nutrients, energy production, metabolism, detoxification and the cycling of neurotransmitters. SNAT3, also designated SN1, G17 and NAT1, is encoded by the human gene SLC38A3. SNAT3 is a glutamine transporter expressed in astroglia from embryonic stages through adulthood, as well as in the liver. Expression levels for SNAT3 in postnatal brain are twice that of normal adult. Increased expression of SNAT3 may also serve as a marker of primary malignant gliomas *in situ*.

REFERENCES

- Wang, H., et al. 2000. Cloning and functional expression of ATA1, a subtype of amino acid transporter A, from human placenta. *Biochem. Biophys. Res. Commun.* 273: 1175-1179.
- Hatanaka, T., et al. 2000. Primary structure, functional characteristics and tissue expression pattern of human ATA2, a subtype of amino acid transport system A. *Biochim. Biophys. Acta* 1467: 1-6.
- Gu, S., et al. 2001. Characterization of an N-system amino acid transporter expressed in retina and its involvement in glutamine transport. *J. Biol. Chem.* 276: 24137-24144.
- Freeman, T.L., et al. 2002. ATA2-mediated amino acid uptake following partial hepatectomy is regulated by redistribution to the plasma membrane. *Arch. Biochem. Biophys.* 400: 215-222.
- Boulland, J.L., et al. 2003. Highly differential expression of SN1, a bidirectional glutamine transporter, in astroglia and endothelium in the developing rat brain. *Glia* 41: 260-275.
- Palii, S.S., et al. 2004. Transcriptional control of the human sodium-coupled neutral amino acid transporter system A gene by amino acid availability is mediated by an intronic element. *J. Biol. Chem.* 279: 3463-3471.

CHROMOSOMAL LOCATION

Genetic locus: SLC38A3 (human) mapping to 3p21.31; Slc38a3 (mouse) mapping to 9 F1.

SOURCE

SNAT3 (N-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of SNAT3 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33445 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS


SNAT3 (N-20) is recommended for detection of SNAT3 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

SNAT3 (N-20) is also recommended for detection of SNAT3 in additional species, including equine, canine, bovine and porcine.


Suitable for use as control antibody for SNAT3 siRNA (h): sc-44980, SNAT3 siRNA (m): sc-44981, SNAT3 shRNA Plasmid (h): sc-44980-SH, SNAT3 shRNA Plasmid (m): sc-44981-SH, SNAT3 shRNA (h) Lentiviral Particles: sc-44980-V and SNAT3 shRNA (m) Lentiviral Particles: sc-44981-V.

Positive Controls: SNAT3 (h): 293T Lysate: sc-115641.

DATA



SNAT3 (N-20): sc-33445. Western blot analysis of SNAT3 expression in non-transfected: sc-117752 (A) and human SNAT3 transfected: sc-115641 (B) 293T whole cell lysates.



SNAT3 (N-20): sc-33445. Immunoperoxidase staining of formalin fixed, paraffin-embedded human liver tissue showing membrane staining of hepatocytes.

SELECT PRODUCT CITATIONS

- Yoshioka, C., et al. 2009. Expression and role of SNAT3 in the placenta. *Placenta* 30: 1071-1077.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **SNAT3 (H-11): sc-398982** or **SNAT3 (H-7): sc-373705**, our highly recommended monoclonal alternatives to SNAT3 (N-20).