

frizzled-9 (D-14): sc-33508

BACKGROUND

The frizzled gene, originally identified in *Drosophila melanogaster*, is involved in the development of tissue polarity. The mammalian homolog of frizzled as well as several secreted mammalian frizzled-related proteins (FRPs) have been described. The frizzled proteins contain seven transmembrane domains, a cysteine-rich domain in the extracellular region and a carboxy terminal Ser/Thr-xxx-Val motif. They function as receptors for Wnt and are generally coupled to G proteins. The frizzled-9 gene is located within the Williams syndrome common deleted region at chromosomal band 7q11.23. Heterozygous deletion of the frizzled-9 gene may contribute to the Williams syndrome phenotype. In mouse, frizzled-9 overexpression can induce the hyperphosphorylation and relocalization of Dvl-1 from the cytoplasm to the cell membrane and cytosolic β -catenin accumulation. In rat, frizzled-9 is important in Wnt/ β -catenin signaling in 293T cells. Frizzled-9 is expressed predominantly in brain, testis, eye, skeletal muscle, and kidney.

REFERENCES

1. Wang, Y., Macke, J.P., Abella, B.S., Andreasson K., Worley P., Gilbert D.J., Copeland, N.G., Jenkins, N.A. and Nathans, J. 1996. A large family of putative transmembrane receptors homologous to the product of the *Drosophila* tissue polarity gene frizzled. *J. Biol. Chem.* 271: 4468-4476.
2. Yang-Snyder, J., Miller, J.R., Brown, J.D., Lai, C.J. and Moon, R.T. 1996. A frizzled homolog functions in a vertebrate Wnt signaling pathway. *Curr. Biol.* 6: 1302-1306.
3. Rattner, A., Hsieh, J.C., Smallwood, P.M., Gilbert, D.J., Copeland, N.G., Jenkins, N.A. and Nathans, J. 1997. A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors. *Proc. Natl. Acad. Sci. USA* 94: 2859-2863.
4. Finch, P.W., He, X., Kelley, M.J., Uren, A., Schaudies, R.P., Popescu, N.C., Rudikoff, S., Aaronson, S.A., Varmus, H.E. and Rubin, J.S. 1997. Purification and molecular cloning of a secreted, frizzled-related antagonist of Wnt action. *Proc. Natl. Acad. Sci. USA* 94: 6770-6775.
5. Melkonyan, H.S., Chang, W.C., Shapiro, J.P., Mahadevappa, M., Fitzpatrick, P.A., Kiefer, M.C., Tomei, L.D. and Umansky, S.R. 1997. SARPs: a family of secreted apoptosis-related proteins. *Proc. Natl. Acad. Sci. USA* 94: 13636-13641.
6. Wang, Y.K., Sporle, R., Paperna, T., Schughart, K. and Francke, U. 1999. Characterization and expression pattern of the frizzled gene Fzd9, the mouse homolog of FZD9 which is deleted in Williams-Beuren syndrome. *Genomics* 57: 235-248.
7. Karasawa, T., Yokokura, H., Kitajewski, J. and Lombroso, P.J. 2002. Frizzled-9 is activated by Wnt-2 and functions in Wnt/ β -catenin signaling. *J. Biol. Chem.* 277: 37479-37486.

CHROMOSOMAL LOCATION

Genetic locus: FZD9 (human) mapping to 7q11.23; Fzd9 (mouse) mapping to 5 G2.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

SOURCE

frizzled-9 (D-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of frizzled-9 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33508 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

frizzled-9 (D-14) is recommended for detection of frizzled-9 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

frizzled-9 (D-14) is also recommended for detection of frizzled-9 in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for frizzled-9 siRNA (h): sc-39994, frizzled-9 siRNA (m): sc-39995, frizzled-9 shRNA Plasmid (h): sc-39994-SH, frizzled-9 shRNA Plasmid (m): sc-39995-SH, frizzled-9 shRNA (h) Lentiviral Particles: sc-39994-V and frizzled-9 shRNA (m) Lentiviral Particles: sc-39995-V.

Molecular Weight of frizzled-9: 65 kDa.

Positive Controls: F9 cell lysate: sc-2245.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.