

frizzled-10 (H-13): sc-33510

BACKGROUND

The frizzled gene, originally identified in *Drosophila melanogaster*, is involved in the development of tissue polarity. The mammalian homolog of frizzled as well as several secreted mammalian frizzled-related proteins (FRPs) have been described. The frizzled proteins contain seven transmembrane domains, a cysteine-rich domain in the extracellular region and a carboxy terminal Ser/Thr-xxx-Val motif. They function as receptors for Wnt and are generally coupled to G proteins. Upregulation of frizzled-10 mRNA in human cells may lead to carcinogenesis through Wnt- β -catenin-TCF signaling pathway activation. Frizzled-10 has been found to be upregulated in HeLa S3, NT2, TMK1 and MKN74 cancer cell lines as well as in colorectal and breast cancer.

REFERENCES

1. Wang, Y., et al. 1996. A large family of putative transmembrane receptors homologous to the product of the *Drosophila* tissue polarity gene frizzled. *J. Biol. Chem.* 271: 4468-4476.
2. Yang-Snyder, J., et al. 1996. A frizzled homolog functions in a vertebrate Wnt signaling pathway. *Curr. Biol.* 6: 1302-1306.
3. Rattner, A., et al. 1997. A family of secreted proteins contains homology to the cysteine-rich ligand-binding domain of frizzled receptors. *Proc. Natl. Acad. Sci. USA* 94: 2859-2863.
4. Finch, P.W., et al. 1997. Purification and molecular cloning of a secreted, Frizzled-related antagonist of Wnt action. *Proc. Natl. Acad. Sci. USA* 94: 6770-6775.
5. Melkonyan, H.S., et al. 1997. SARPs: a family of secreted apoptosis-related proteins. *Proc. Natl. Acad. Sci. USA* 94: 13636-13641.

CHROMOSOMAL LOCATION

Genetic locus: FZD10 (human) mapping to 12q24.33, FZD9 (human) mapping to 7q11.23; Fzd10 (mouse) mapping to 5 G1.3, Fzd9 (mouse) mapping to 5 G2.

SOURCE

frizzled-10 (H-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of frizzled-10 of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33510 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

frizzled-10 (H-13) is recommended for detection of frizzled-10 and frizzled-9 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

frizzled-10 (H-13) is also recommended for detection of frizzled-10 and frizzled-9 in additional species, including equine, canine, bovine, porcine and avian.


Molecular Weight of frizzled-10: 73 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200 or rat postnatal kidney tissue extract.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



frizzled-10 (H-13): sc-33510. Western blot analysis of frizzled-10 expression in rat postnatal kidney tissue extract.

RESEARCH USE

For research use only, not for use in diagnostic procedures.