

δ-catenin (H-160): sc-33553

BACKGROUND

The catenins, α , β , γ and δ , are proteins which bind to the highly conserved, intracellular cytoplasmic tail of E-cadherin. Together, the catenin/cadherin complexes play an important role mediating cellular adhesion. α -catenin was initially described as an E-cadherin associated protein, and since has been shown to associate with other members of the cadherin family, such as N-cadherin and P-cadherin. β -catenin associates with the cytoplasmic portion of E-cadherin, which is necessary for the function of E-cadherin as an adhesion molecule. β -catenin has also been found in complexes with the tumor suppressor protein APC. δ -catenin, also known as plakoglobin, binds with α -catenin and N-cadherin. δ -catenin interacts with presenilin 1 and is expressed in the brain. The gene encoding δ -catenin maps to human chromosome 5p15.2. A hemizygous loss of the gene encoding δ -catenin leads to the mental retardation associated with Cri-du-Chat syndrome. In addition, the transmembrane phosphatase PTPm associates with catenin/cadherin complexes and may regulate complex signaling.

CHROMOSOMAL LOCATION

Genetic locus: CTNND2 (human) mapping to 5p15.2; Ctnnd2 (mouse) mapping to 15 B2.

SOURCE

δ -catenin (H-160) is a rabbit polyclonal antibody raised against amino acids 1031-1190 mapping near the C-terminus of δ -catenin of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

δ -catenin (H-160) is recommended for detection of δ -catenin of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

δ -catenin (H-160) is also recommended for detection of δ -catenin in additional species, including equine, canine and bovine.

Suitable for use as control antibody for δ -catenin siRNA (h): sc-43021, δ -catenin siRNA (m): sc-43022, δ -catenin shRNA Plasmid (h): sc-43021-SH, δ -catenin shRNA Plasmid (m): sc-43022-SH, δ -catenin shRNA (h) Lentiviral Particles: sc-43021-V and δ -catenin shRNA (m) Lentiviral Particles: sc-43022-V.


Molecular Weight of δ -catenin: 133 kDa.

Positive Controls: PC-12 cell lysate: sc-2250, HeLa whole cell lysate: sc-2200 or A-431 whole cell lysate: sc-2201.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



δ -catenin (H-160): sc-33553. Western blot analysis of δ -catenin expression in PC-12 whole cell lysate.



δ -catenin (H-160): sc-33553. Immunofluorescence staining of formalin-fixed Hep G2 cells showing cytoplasmic localization.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **δ -catenin (40.1): sc-81793**, our highly recommended monoclonal alternative to δ -catenin (H-160).