

Glycophorin B (H-46): sc-33561

BACKGROUND

Glycophorins A, B and C are sialoglycoproteins of the human erythrocyte membrane which bear the antigenic determinants for the MN, Ss and Gerbich blood groups, respectively. Glycophorins span the membrane once and present their amino-terminal end to the extracellular surface of the human erythrocyte. The genetic array of expressed Glycophorin surface antigens on erythrocytes defines the blood group phenotype of the individual. The human Glycophorin A gene maps to chromosome 4q31.22, contains seven exons which are 97% homologous to Glycophorin B and encodes a 150 amino acid protein. The human Glycophorin B gene maps to chromosome 4q31.22 and encodes a 91 amino acid protein. The human Glycophorin C gene maps to chromosome 2q14.3 and contains four exons. Glycophorin C transcript can generate two protein isoforms. Isoform 1 includes all 4 exons and encodes the full length, 128 amino acid protein. Isoform 2 is missing exon 2 and encodes a 109 amino acid protein, which specifies the Yus subtype of the Gerbich phenotype.

REFERENCES

1. Chang, S.H., et al. 2001. Regulation of the Glycophorin C protein 4.1 membrane-to-skeleton bridge and evaluation of its contribution to erythrocyte membrane stability. *J. Biol. Chem.* 276: 22223-22230.
2. Gerber, D., et al. 2001. *In vivo* detection of hetero-association of Glycophorin A and its mutants within the membrane. *J. Biol. Chem.* 276: 31229-31232.
3. Lobo, C.A., et al. 2003. Glycophorin C is the receptor for the *Plasmodium falciparum* erythrocyte binding ligand PfEBP-2 (baebl). *Blood* 101: 4628-4631.
4. Young, M.T., et al. 2003. Distinct regions of human Glycophorin A enhance human red cell anion exchanger (band 3; AE1) transport function and surface trafficking. *J. Biol. Chem.* 278: 32954-32961.
5. Overton, M.C., et al. 2003. Oligomerization, biogenesis and signaling is promoted by a Glycophorin A-like dimerization motif in transmembrane domain 1 of a yeast G protein-coupled receptor. *J. Biol. Chem.* 278: 49369-49377.
6. Bruce, L.J., et al. 2004. Altered structure and anion transport properties of band 3 (AE1, SLC4A1) in human red cells lacking Glycophorin A. *J. Biol. Chem.* 279: 2414-2420.
7. Lahlil, R., et al. 2004. SCL assembles a multifactorial complex that determines Glycophorin A expression. *Mol. Cell. Biol.* 24: 1439-1452.
8. Gerber, D., et al. 2004. Structural adaptation of the Glycophorin A transmembrane homodimer to D-amino acid modifications. *J. Mol. Biol.* 339: 243-250.

CHROMOSOMAL LOCATION

Genetic locus: GYPB (human) mapping to 4q31.22.

SOURCE

Glycophorin B (H-46) is a rabbit polyclonal antibody raised against amino acids 46-91 mapping at the C-terminus of Glycophorin B of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Glycophorin B (H-46) is recommended for detection of Glycophorin B, and to a lesser extent, Glycophorin A of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of Glycophorin B (monomer): 25 kDa.


Molecular Weight of Glycophorin B (dimer): 46 kDa.

Positive Controls: human spleen tissue.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



Glycophorin B (H-46): sc-33561. Western blot analysis of human recombinant Glycophorin B fusion protein.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.