

OATP-C (H-60): sc-33609

BACKGROUND

The organic anion transporting polypeptides, OATP-A (also designated OATP1, OATP1A2 and SLC21A3) and OATP-C (also designated OATP2, SLC21A6 and LST1), mediate hepatic uptake of cardiac glycosides. The expression of OATP-C, but not OATP-A, is inducible by phenobarbital and pregnenolone-16 α -carbonitrile, resulting in the increased capacity of the liver to extract cardiac glycosides from the plasma. OATP-A is expressed in liver and kidney and helps mediate sodium-independent uptake of the anionic steroid conjugates dehydroepiandrosterone sulfate, estradiol-17-glucuronide and prostaglandin. OATP-C is exclusively expressed in liver and is localized to the basolateral hepatocyte membrane. Although OATP-C mRNA levels decrease during pregnancy and increase postpartum, OATP-C protein levels remain relatively constant. OATP-C transports taurocholic acid, the adrenal androgen dehydroepiandrosterone sulfate, thyroid hormone, hydroxymethylglutaryl-CoA reductase inhibitor and pravastatin. OATP-C is therefore a novel organic anion transport protein that has overlapping but not identical substrate specificities with other subtypes of OATP. OATP-A and OATP-C are both pravastatin transporters, suggesting that they are responsible for the hepatic uptake of the liver-specific hydroxymethylglutaryl-CoA reductase inhibitor in mouse, rat and human.

REFERENCES

- Hsiang, B., et al. 1999. A novel human hepatic organic anion transporting polypeptide (OATP2). *J. Biol. Chem.* 274: 37161-37168.
- König, J., et al. 2000. Localization and genomic organization of a new hepatocellular organic anion transporting polypeptide. *J. Biol. Chem.* 275: 23161-23168.
- Cattori, V., et al. 2000. Identification of organic anion transporting polypeptide 4 (Oatp4) as a major full-length isoform of the liver-specific transporter-1 (rlst-1) in rat liver. *FEBS Letts.* 474: 242-245.
- König, J., et al. 2000. A novel human organic anion transporting polypeptide localized to the basolateral hepatocyte membrane. *Am. J. Physiol. Gastrointest. Liver Physiol.* 278: 156-164.
- Cao, J., et al. 2001. Differential regulation of hepatic bile salt and organic anion transporters in pregnant and postpartum rats and the role of prolactin. *Hepatology* 33: 140-147.

CHROMOSOMAL LOCATION

Genetic locus: SLC01B1 (human) mapping to 12p.2.

SOURCE

OATP-C (H-60) is a rabbit polyclonal antibody raised against amino acids 606-665 mapping near the C-terminus of OATP-C of human origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

OATP-C (H-60) is recommended for detection of OATP-C and, to a lesser extent, OATP8 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for OATP-C siRNA (h): sc-42549, OATP-C shRNA Plasmid (h): sc-42549-SH and OATP-C shRNA (h) Lentiviral Particles: sc-42549-V.

Molecular Weight of OATP-C: 90 kDa.

Positive Controls: SW480 cell lysate:0 sc-2219 or HeLa whole cell lysate: sc-2200.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.


 MONOS
 Satisfation
 Guaranteed

Try **OATP-C (A-3): sc-271157**, our highly recommended monoclonal alternative to OATP-C (H-60).