

# EPI64 (H-128): sc-33616

## BACKGROUND

The Na<sup>+</sup>/H<sup>+</sup> exchange protein (NHE) functions in transepithelial Na<sup>+</sup> absorption and is primarily expressed in the intestinal and renal brush border membrane. NHE regulatory factor 1 (NHERF-1) interacts with NHE through two PDZ (for PSD-95, discs-large and ZO-1 homology) domains, which are protein-protein interaction modules that associate with specific C-terminal motifs on target proteins. Also known as EBP50, NHERF-1 facilitates cAMP inhibition of NHE to decrease Na<sup>+</sup> adsorption. NHERF-2, also known as E3KARP, is ubiquitously expressed as a 34 kDa protein which also functions in NHE2 regulation. EBP-PDZ interactor of 64 kDa (EPI64) contains a C-terminal-DTYL sequence that binds to the first PDZ domain of NHERF-1 and NHERF-2. EPI64 is ubiquitously expressed and localizes with NHERF-1 *in vitro*. The gene encoding human EPI64 maps to chromosome 22q12.1.

## REFERENCES

- Sheng, M. 1996. PDZs and receptor/channel clustering: rounding up the latest suspects. *Neuron* 17: 575-578.
- Yun, C.H., Oh, S., Zizak, M., Steplock, D., Tsao, S., Tse, C.M., Weinman, E.J. and Donowitz, M. 1997. cAMP-mediated inhibition of the epithelial brush border Na<sup>+</sup>/H<sup>+</sup> exchanger, NHE3, requires an associated regulatory protein. *Proc. Natl. Acad. Sci. USA* 94: 3010-3015.
- Poulat, F., de Santa Barbara, P., Desclozeaux, M., Soullier, S., Moniot, B., Bonneaud, N., Boizet, B. and Berta, P. 1997. The human testis determining factor SRY bind a nuclear factor containing PDZ protein interaction domains. *J. Biol. Chem.* 272: 7167-7172.
- Reczek, D. and Bretscher, A. 2001. Identification of EPI64, a TBC/rabGAP domain-containing microvillar protein that binds to the first PDZ domain of EBP50 and E3KARP. *J. Cell Biol.* 153: 191-205.
- Itoh, T and Fukuda, M. 2006. Identification of EPI64 as a GTPase-activating protein specific for Rab27A. *J. Biol. Chem.* 281: 31823-31831.
- Hanono, A., Garbett, D., Reczek, D., Chambers, D.N. and Bretscher, A. 2006. EPI64 regulates microvillar subdomains and structure. *J. Cell Biol.* 175: 803-813.

## CHROMOSOMAL LOCATION

Genetic locus: TBC1D10A (human) mapping to 22q12.1; Tbc1d10a (mouse) mapping to 11 A1.

## SOURCE

EPI64 (H-128) is a rabbit polyclonal antibody raised against amino acids 381-508 mapping at the C-terminus of EPI64 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

EPI64 (H-128) is recommended for detection of EPI64 of human and, to a lesser extent, mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for EPI64 siRNA (h): sc-42520, EPI64 siRNA (m): sc-42521, EPI64 shRNA Plasmid (h): sc-42520-SH, EPI64 shRNA Plasmid (m): sc-42521-SH, EPI64 shRNA (h) Lentiviral Particles: sc-42520-V and EPI64 shRNA (m) Lentiviral Particles: sc-42521-V.


Molecular Weight of EPI64: 56 kDa.

Positive Controls: EPI64 (h3): 293T Lysate: sc-128540 or mouse embryo tissue extract.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



EPI64 (H-128): sc-33616. Western blot analysis of EPI64 expression in non-transfected: sc-117752 (A) and human EPI64 transfected: sc-128540 (B) 293T whole cell lysates.



EPI64 (H-128): sc-33616. Western blot analysis of EPI64 expression in mouse embryo tissue extract.

## RESEARCH USE

For research use only, not for use in diagnostic procedure.

**MONOS**  
Satisfaction  
Guaranteed

Try **EPI64 (34): sc-136520** or **EPI64 (B-9): sc-376991**, our highly recommended monoclonal alternatives to EPI64 (H-128).