

Cas-L (2B11): sc-33658

BACKGROUND

Cas family proteins are adhesion docking molecules that mediate protein-protein interactions and contribute to a number of signal transduction pathways. Cas-L (also designated human enhancer of filamentation (HEF1) and neural precursor cell expressed, developmentally down-regulated 9 (NEDD9), participates in integrin and growth factor signaling pathways that regulate growth, motility and apoptosis. Cas-L consists of two isoforms, p105 and p115. The larger molecular weight form is a result of Ser/Thr phosphorylation. Cas-L phosphorylation is dependent on cell adhesion and Src kinase activity. Cas-L acts as a downstream effector of FAK in the invasive behavior of glioblastoma cells. TGF β 1 regulates Cas-L gene expression and influences phosphorylation. Adhesion-dependent Actin organization regulates proteasomal turnover of Cas-L through the activity of PP2A. Tyrosine phosphorylated Cas-L can bind FAK in dendrite and soma of neurons after ischemia. Cas-L can promote neurite outgrowth of PC-12 cells.

REFERENCES

- Liu, X., et al. 2000. A novel ability of Smad3 to regulate proteasomal degradation of a Cas family member HEF1. *EMBO J.* 19: 6759-6769.
- Law, S.F., et al. 2000. The docking protein HEF1 is an apoptotic mediator at focal adhesion sites. *Mol. Cell. Biol.* 20: 5184-5195.
- Zheng, M., et al. 2002. Regulation of HEF1 expression and phosphorylation by TGF β -1 and cell adhesion. *J. Biol. Chem.* 277: 39599-39608.
- Pugacheva, E.N., et al. 2005. The focal adhesion scaffolding protein HEF1 regulates activation of the Aurora-A and Nek2 kinases at the centrosome. *Nat. Cell Biol.* 7: 937-946.
- Iwata, S., et al. 2005. HTLV-I Tax induces and associates with Crk-associated substrate lymphocyte type (Cas-L). *Oncogene* 24: 1262-1271.
- Sasaki, T., et al. 2005. Nedd9 protein, a Cas-L homologue, is upregulated after transient global ischemia in rats: possible involvement of Nedd9 in the differentiation of neurons after ischemia. *Stroke* 36: 2457-2462.
- Natarajan, M., et al. 2006. HEF1 is a necessary and specific downstream effector of FAK that promotes the migration of glioblastoma cells. *Oncogene* 25: 1721-1732.
- Zheng, M., et al. 2006. Cell adhesion regulates Ser/Thr phosphorylation and proteasomal degradation of HEF1. *J. Cell Sci.* 119: 96-103.

CHROMOSOMAL LOCATION

Genetic locus: NEDD9 (human) mapping to 6p24.2; Nedd9 (mouse) mapping to 13 A3.3.

SOURCE

Cas-L (2B11) is a mouse monoclonal antibody raised against amino acids 82-398 of Cas-L of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Cas-L (2B11) is recommended for detection of Cas-L of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for Cas-L siRNA (h): sc-40794, Cas-L siRNA (m): sc-40795, Cas-L shRNA Plasmid (h): sc-40794-SH, Cas-L shRNA Plasmid (m): sc-40795-SH, Cas-L shRNA (h) Lentiviral Particles: sc-40794-V and Cas-L shRNA (m) Lentiviral Particles: sc-40795-V.


Molecular Weight of Cas-L: 105 kDa.

Positive Controls: MCF7 whole cell lysate: sc-2206, A549 cell lysate: sc-2413 or Cas-L (h2): 293T Lysate: sc-159485.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

DATA



Cas-L (2B11): sc-33658. Western blot analysis of Cas-L expression in MCF7 (A), A549 (B), NIH/3T3 (C) and A-10 (D) whole cell lysates.



Cas-L (2B11): sc-33658. Western blot analysis of Cas-L expression in non-transfected: sc-117752 (A) and human Cas-L transfected: sc-159485 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Xia, D., et al. 2010. HEF1 is a crucial mediator of the proliferative effects of prostaglandin E $_2$ on colon cancer cells. *Cancer Res.* 70: 824-831.

STORAGE

Store at 4 $^{\circ}$ C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.