

neurocan (650.24): sc-33663

BACKGROUND

The lecticans are a family of chondroitin proteoglycans, including aggrecan, versican, neurocan and brevican, that contain a C-type lectin domain. Neurocan is a glycoprotein synthesized primarily by neurons, and its expression levels are highest during embryonic brain development and during the early postnatal period. Neurocan is a component of the extracellular matrix in the central nervous system that can bind to various other CNS matrix components, such as heparin, Tenascin-R, HB-GAM and NCAM, suggesting that it plays a role in axon guidance and neurite growth. Neurocan is a 1,257 amino acid, precursor protein in embryonic and neonatal rats that is proteolytically processed in the adult CNS into an N-terminal fragment, which localizes to the cytoplasm of glial cells. Neurocan expression (including the re-expression of the neonatal isoform) increases significantly in adults in the glial scar resulting from cortical injury.

REFERENCES

1. Retzler, C., et al. 1996. Structural and electron microscopic analysis of neurocan and recombinant neurocan fragments. *J. Biol. Chem.* 271: 17107-17113.
2. Milev, P., et al. 1998. High affinity binding and overlapping localization of neurocan and phosphacan/protein-tyrosine phosphatase- ζ/β with Tenascin-R, amphoterin and the heparin-binding growth associated molecule. *J. Biol. Chem.* 273: 6998-7005.
3. Milev, P., et al. 1998. Differential regulation of expression of hyaluronan-binding proteoglycans in developing brain: aggrecan, versican, neurocan and brevican. *Biochem. Biophys. Res. Commun.* 247: 207-212.
4. Matsui, F., et al. 1998. Occurrence of a N-terminal proteolytic fragment of neurocan, not a C-terminal half, in perineuronal net in the adult rat cerebrum. *Brain Res.* 790: 45-51.

CHROMOSOMAL LOCATION

Genetic locus: NCAN (human) mapping to 19p13.11; Ncan (mouse) mapping to 8 B3.3.

SOURCE

neurocan (650.24) is a mouse monoclonal antibody immunized with purified proteoglycan fraction of neonatal rat brain.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

neurocan (650.24) is available conjugated to agarose (sc-33663 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-33663 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-33663 PE), fluorescein (sc-33663 FITC), Alexa Fluor[®] 488 (sc-33663 AF488), Alexa Fluor[®] 546 (sc-33663 AF546), Alexa Fluor[®] 594 (sc-33663 AF594) or Alexa Fluor[®] 647 (sc-33663 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-33663 AF680) or Alexa Fluor[®] 790 (sc-33663 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

neurocan (650.24) is recommended for detection of neurocan of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).


Suitable for use as control antibody for neurocan siRNA (h): sc-41901, neurocan siRNA (m): sc-41902, neurocan shRNA Plasmid (h): sc-41901-SH, neurocan shRNA Plasmid (m): sc-41902-SH, neurocan shRNA (h) Lentiviral Particles: sc-41901-V and neurocan shRNA (m) Lentiviral Particles: sc-41902-V.

Molecular Weight of full length neurocan: 220 kDa.


Molecular Weight of neurocan proteolytic variant: 140 kDa.

Positive Controls: rat brain extract: sc-2392, rat postnatal brain tissue extract or mouse brain extract: sc-2253.

DATA



neurocan (650.24): sc-33663. Western blot analysis of neurocan expression in mouse brain tissue extract. Detection reagent used: m-IgG κ BP-HRP: sc-516102.



neurocan (650.24): sc-33663. Western blot analysis of neurocan expression in rat brain tissue extract. Detection reagent used: m-IgG κ BP-HRP: sc-516102.

SELECT PRODUCT CITATIONS

1. Gao, Q.S., et al. 2021. Brief inhalation of sevoflurane can reduce glial scar formation after hypoxic-ischemic brain injury in neonatal rats. *Neural Regen Res.* 16: 1052-1061.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.