



## HID (d-300): sc-33744

### BACKGROUND

*Drosophila melanogaster* is a proven and effective model for studying developmental and cellular processes common to higher eukaryotes. Approximately 13,600 genes have been elucidated from more than 120 megabases of euchromatin, and they are organized among the chromosomes 2, 3, 4, X and Y, with the Y chromosome being predominately heterochromatic. *Drosophila* genes can be categorized based on the type of protein for which they encode and are represented by six major classifications, which include intracellular signaling proteins, transmembrane proteins, RNA binding proteins, secreted factors, transcription regulators (basic helix-loop-helix, homeodomain containing, zinc finger containing, and chromatin associated) or other functional proteins. Among these numerous proteins, HID (head involution defective, WRINKLED) is a 410 amino acid, death domain-containing proapoptotic protein that contributes to proper head and wing development.

### REFERENCES

- Grether, M.E., Abrams, J.M., Agapite, J., White, K. and Steller, H. 1995. The head involution defective gene of *Drosophila melanogaster* functions in programmed cell death. *Genes Dev.* 9: 1694-1708.
- Bergmann, A., Agapite, J., McCall, K. and Steller, H. 1998. The *Drosophila* gene HID is a direct molecular target of Ras-dependent survival signaling. *Cell* 95: 331-341.
- Adams, M.D., Celniker, S.E., Holt, R.A., Evans, C.A., Gocayne, J.D. and Amanatides, P., et al. 2000. The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185-2195.
- Goyal, L., McCall, K., Agapite, J., Hartwig, E. and Steller, H. 2000. Induction of apoptosis by *Drosophila* reaper, HID and GRIM through inhibition of IAP function. *EMBO J.* 19: 589-597.
- The Interactive Fly. <http://www.sdbonline.org/fly/aimain/1aahome.htm>  
<http://www.sdbonline.org/fly/dbzhnsky/headind1.htm>
- LocusLink Report (LocusID: 40009). <http://www.ncbi.nlm.nih.gov/LocusLink/>

### SOURCE

HID (d-300) is a rabbit polyclonal antibody raised against amino acids 1-300 mapping at the N-terminus of HID of *Drosophila melanogaster* origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

### STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


### APPLICATIONS

HID (d-300) is recommended for detection of HID of *Drosophila melanogaster* origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1–2 µg per 100–500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### DATA



HID (d-300): sc-33744. Western blot analysis of *Drosophila* recombinant HID fusion protein.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.