

# Gonadotropin $\alpha$ (FL-116): sc-33793

## BACKGROUND

Gonadotropin (also designated choriogonadotropin) is a hormone produced by the placenta in the first trimester of pregnancy and exists as a hetero-dimer formed from a common  $\alpha$  chain and a unique  $\beta$  chain. The unique  $\beta$  chain confers biological specificity to thyrotropin, lutropin, follitropin and gonadotropin. The secreted  $\alpha$  subunit maps to human chromosome 6 and the  $\beta$  subunit maps to human chromosome 19. Gonadotropin stimulates the ovaries to produce and maintain normal levels of the steroids essential for maintaining pregnancy, including estrogen and progesterone. Gonadotropin is a member of the cystine knot growth-factor superfamily, a group of proteins that contain a distinct arrangement of six cysteine residues and are expressed in placenta. The proper secretion and dimerization of gonadotropin depends on the conformation of the cystine knot, although biological activity is independent of this conformation.

## REFERENCES

- Naylor, S.L., et al. 1983. Chromosome assignment of the genes encoding the  $\alpha$  and  $\beta$  subunits of the glycoprotein hormones in man and mouse. *Somatic Cell Genet.* 9: 757-770.
- Laphorn, A.J., et al. 1994. Crystal structure of human chorionic gonadotropin. *Nature* 369: 455-461.
- Furuhashi, M., et al. 1994. Mutagenesis of cysteine residues in the human gonadotropin  $\alpha$  subunit. Roles of individual disulfide bonds in secretion, assembly, and biologic activity. *J. Biol. Chem.* 269: 25543-25548.
- Sun, P.D., et al. 1995. The cystine-knot growth-factor superfamily. *Annu. Rev. Biophys. Biomol. Struct.* 24: 269-291.
- Furuhashi, M., et al. 1996. Disulfide bonds 7-31 and 59-87 of the  $\alpha$  subunit play a different role in assembly of human chorionic gonadotropin and lutropin. *Endocrinology* 137: 4196-4200.
- Sato, A., et al. 1997. Cystine knot of the gonadotropin  $\alpha$  subunit is critical for intracellular behavior but not for *in vitro* biological activity. *J. Biol. Chem.* 272: 18098-18103.
- Lustbader, J.W., et al. 1998. Structural and molecular studies of human chorionic gonadotropin and its receptor. *Recent Prog. Horm. Res.* 53: 395-424.
- Vaananen, J.E., et al. 1998. Regulation of prostaglandin F $2\alpha$ -receptor mRNA in human granulosa-luteal cells by human chorionic gonadotropin and prostaglandin. *Endocrine* 8: 261-267.

## CHROMOSOMAL LOCATION

Genetic locus: CGA (human) mapping to 6q14.3.

## SOURCE

Gonadotropin  $\alpha$  (FL-116) is a rabbit polyclonal antibody raised against amino acids 57-116 mapping at the C-terminus of Gonadotropin  $\alpha$  of human origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

Gonadotropin  $\alpha$  (FL-116) is recommended for detection of precursor and mature Gonadotropin  $\alpha$  of human and, to a lesser extent, rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Gonadotropin  $\alpha$  siRNA (h): sc-39538, Gonadotropin  $\alpha$  shRNA Plasmid (h): sc-39538-SH and Gonadotropin  $\alpha$  shRNA (h) Lentiviral Particles: sc-39538-V.


Molecular Weight of Gonadotropin  $\alpha$ : 23.5 kDa.

Positive Controls: rat pituitary gland extract: sc-364807.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Gonadotropin  $\alpha$  (FL-116): sc-33793. Western blot analysis of Gonadotropin  $\alpha$  expression in rat pituitary tissue extract.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.