

# CRISP-1 (H-75): sc-33802

## BACKGROUND

Cysteine-rich secretory proteins (CRISPs) represent a family of evolutionarily conserved proteins which may play a role in the innate immune system and are transcriptionally regulated by androgens in several tissues. AEG is a sperm surface protein involved in the fusion of egg and sperm. Although CRISP-1 (also designated AEG-like protein, ARP, cysteine-rich secretory protein-1 or AEG-related protein) is not the ortholog of rodent AEG, it resembles AEG in that it is an epididymal secretory glycoprotein that binds to the postacrosomal region of the sperm head. CRISP-1 coats the postacrosomal region of sperm heads as they pass through the epididymis. CRISP-1 is found in all regions of the epididymis, ductus deferens, seminal plasma and sperm. CRISP-3 is expressed in pancreas and prostate tissues, and along with CRISP-1, is expressed in saliva. The gene that encodes CRISP-3 is an early response gene that may participate in the pathophysiology of the autoimmune lesions of Sjogren's syndrome.

## REFERENCES

- Hayashi, M., Fujimoto, S., Takano, H., Ushiki, T., Abe, K., Ishikura, H., Yoshida, M.C., Kirchhoff, C., Ishibashi, T. and Kasahara, M. 1996. Characterization of a human glycoprotein with a potential role in sperm-egg fusion: cDNA cloning, immunohistochemical localization, and chromosomal assignment of the gene (AEG1). *Genomics* 32: 367-374.
- Schwidetzky, U., Schleuning, W.D. and Haendler, B. 1997. Isolation and characterization of the androgen-dependent mouse cysteine-rich secretory protein-1 (CRISP-1) gene. *Biochem. J.* 321: 325-332.
- Haendler, B., Habenicht, U.F., Schwidetzky, U., Schuttke, I. and Schleuning, W.D. 1997. Differential androgen regulation of the murine genes for cysteine-rich secretory proteins (CRISP). *Eur. J. Biochem.* 250: 440-446.
- Haendler, B., Toda, I., Sullivan, D.A. and Schleuning, W.D. 1999. Expression of transcripts for cysteine-rich secretory proteins (CRISPs) in the murine lacrimal gland. *J. Cell. Physiol.* 178: 371-378.
- Tapinos, N.I., Polihronis, M., Thyphronitis, G. and Moutsopoulos, H.M. 2002. Characterization of the cysteine-rich secretory protein 3 gene as an early-transcribed gene with a putative role in the pathophysiology of Sjogren's syndrome. *Arthritis Rheum.* 46: 215-222.

## CHROMOSOMAL LOCATION

Genetic locus: CRISP1 (human) mapping to 6p12.3.

## SOURCE

CRISP-1 (H-75) is a rabbit polyclonal antibody raised against amino acids 66-140 mapping within an internal region of CRISP-1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

CRISP-1 (H-75) is recommended for detection of CRISP-1 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for CRISP-1 siRNA (h): sc-44749, CRISP-1 shRNA Plasmid (h): sc-44749-SH and CRISP-1 shRNA (h) Lentiviral Particles: sc-44749-V.


Molecular Weight of CRISP-1: 30 kDa.

Positive Controls: Hep G2 cell lysate: sc-2227 or human testis extract: sc-363781.

## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



CRISP-1 (H-75): sc-33802. Western blot analysis of CRISP-1 expression in Hep G2 whole cell lysate (A) and human testis tissue extract (B).

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.