

Apaf-1 (D-18): sc-33870

BACKGROUND

The mammalian homologs of the Ced-4 proteins, Apaf-1 (Ced-4), Nod1 (CARD4), and Nod2 contain a caspase recruitment domain (CARD) and a putative nucleotide binding domain, signified by a consensus Walker's A box (P-loop) and B box (Mg²⁺-binding site). Nod1 contains a putative regulatory domain and multiple leucine-rich repeats. Nod1 is a member of a growing family of intracellular proteins which share structural homology to the apoptosis regulator Apaf-1. Nod1 associates with the CARD-containing kinase RICK and activates NFκB. The self-association of Nod1 mediates proximity of RICK and the interaction of RICK with IKKγ. In addition, Nod-1 binds to multiple caspases with long prodomains, but specifically activates caspase-9 and promotes caspase-9-induced apoptosis. Nod2 is composed of two N-terminal CARDS, a nucleotide-binding domain, and multiple C-terminal leucine-rich repeats. The expression of Nod2 is highly restricted to monocytes, and activates NFκB in response to bacterial lipopolysaccharides.

CHROMOSOMAL LOCATION

Genetic locus: APAF1 (human) mapping to 12q23.1; Apaf1 (mouse) mapping to 10 C2.

SOURCE

Apaf-1 (D-18) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of Apaf-1 of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-33870 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

Apaf-1 (D-18) is recommended for detection of Apaf-1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000). Apaf-1 (D-18) is also recommended for detection of Apaf-1 in additional species, including bovine.

Suitable for use as control antibody for Apaf-1 siRNA (h): sc-29201, Apaf-1 siRNA (m): sc-37147, Apaf-1 shRNA Plasmid (h): sc-29201-SH, Apaf-1 shRNA Plasmid (m): sc-37147-SH, Apaf-1 shRNA (h) Lentiviral Particles: sc-29201-V and Apaf-1 shRNA (m) Lentiviral Particles: sc-37147-V.


Molecular Weight of Apaf-1: 130 kDa.

Positive Controls: WI-38 whole cell lysate: sc-364260, ECV304 cell lysate: sc-2269 or HUV-EC-C whole cell lysate: sc-364180.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



Apaf-1 (D-18): sc-33870. Western blot analysis of Apaf-1 expression in HUV-EC-C (A), ECV304 (B), K-562 (C), human PBL (D) and WI 38 (E) whole cell lysates.



Apaf-1 (D-18): sc-33870. Immunoperoxidase staining of formalin fixed, paraffin-embedded human spleen tissue showing cytoplasmic staining of cells in white pulp and cells in red pulp.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **Apaf-1 (18H2): sc-135624** or **Apaf-1 (24): sc-135836**, our highly recommended monoclonal alternatives to Apaf-1 (D-18).