

# AQP1 (T-13): sc-34008

## BACKGROUND

Aquaporins (AQPs) are a large family of integral membrane water transport channel proteins that facilitate the transport of water through the cell membrane. This function is conserved in animals, plants and bacteria. Many isoforms of Aquaporin have been identified in mammals, designated AQP0 through AQP10. Aquaporins are widely distributed and it is not uncommon for more than one type of AQP to be present in the same cell. Although most Aquaporins are only permeable to water, AQP3, AQP7, AQP9 and one of the two AQP10 transcripts are also permeable to urea and glycerol. AQP2 is the only water channel that is activated by vasopressin to enhance water reabsorption in the kidney collecting duct. Aquaporins are involved in renal water absorption, generation of pulmonary secretions, lacrimation and the secretion and reabsorption of cerebrospinal fluid and aqueous humor. AQP1 is an integral membrane protein expressed in erythrocytes and renal tubule cells.

## CHROMOSOMAL LOCATION

Genetic locus: AQP1 (human) mapping to 7p14.3; Aqp1 (mouse) mapping to 6 B3.

## SOURCE

AQP1 (T-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of AQP1 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34008 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## APPLICATIONS

AQP1 (T-13) is recommended for detection of AQP1 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for AQP1 siRNA (h): sc-29711, AQP1 siRNA (m): sc-29712, AQP1 siRNA (r): sc-156108, AQP1 shRNA Plasmid (h): sc-29711-SH, AQP1 shRNA Plasmid (m): sc-29712-SH, AQP1 shRNA Plasmid (r): sc-156108-SH, AQP1 shRNA (h) Lentiviral Particles: sc-29711-V, AQP1 shRNA (m) Lentiviral Particles: sc-29712-V and AQP1 shRNA (r) Lentiviral Particles: sc-156108-V.

Molecular Weight of AQP1: 28 kDa.


Molecular Weight of glycosylated AQP1: 35-45 kDa.

Positive Controls: Human kidney extract: sc-363764, mouse kidney extract: sc-2255 or rat kidney extract: sc-2394.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 4) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

## DATA



AQP1 (T-13): sc-34008. Western blot analysis of AQP1 expression in human kidney tissue extract.



AQP1 (T-13): sc-34008. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing membrane staining of centroacinar cells.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **AQP1 (B-11): sc-25287** or **AQP1 (1/22): sc-32737**, our highly recommended monoclonal alternatives to AQP1 (T-13). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **AQP1 (B-11): sc-25287**.