

# Ac-Histone H4 (Ser 1/Lys 5/Lys 8/Lys 12): sc-34263

## BACKGROUND

In eukaryotes, DNA is wrapped around histone octamers to form the basic unit of chromatin structure. The octamer is composed of Histones H2A, H2B, H3 and H4, and it associates with approximately 200 base pairs of DNA to form the nucleosome. The association of DNA with histones results in dense packing of chromatin, which restricts proteins involved in gene transcription from binding to DNA. p300 preferentially acetylates Histone H3 at Lysines 14 and 18 and Histone H4 at Lysines 5 and 8. PCAF in its native form primarily acetylates Histone H3 at Lysine 14 to a monoacetylated form and less efficiently acetylates Histone H4 at Lysine 8. Histone H4 may also be acetylated at Lysines 12 and 16, and the involvement of acetylated H4 with Histones H2A, H2B and H3 suggests that acetylated Histones may be involved in dynamic chromatin remodeling.

## SOURCE

Ac-Histone H4 (Ser 1/Lys 5/Lys 8/Lys 12) is a rabbit polyclonal antibody raised against a short amino acid sequence containing acetylated serine 1 and lysine 5, 8 and 12 of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34263 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Ac-Histone H4 (Ser 1/Lys 5/Lys 8/Lys 12) is recommended for detection of Serine 1 and Lysine 5, 8 and 12 acetylated Histone H4 of broad species origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); non cross-reactive with non-acetylated Histone H4 or other lysine acetylation sites.

Ac-Histone H4 (Ser 1/Lys 5/Lys 8/Lys 12) is also recommended for detection of Serine 1 and Lysine 5, 8 and 12 acetylated Histone H4 in additional species, including equine, canine, bovine, porcine and avian.

Molecular Weight of acetylated Ac-Histone H4: 11 kDa.

Molecular Weight of non-acetylated Ac-Histone H4: 11 kDa.


Molecular Weight of hyper-acetylated Ac-Histone H4: 35 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, HeLa nuclear extract: sc-2120 or IMR-32 nuclear extract: sc-2148.


## RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-rabbit IgG-HRP: sc-2004 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible goat anti-rabbit IgG-HRP: sc-2030 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use goat anti-rabbit IgG-FITC: sc-2012 (dilution range: 1:100-1:400) or goat anti-rabbit IgG-TR: sc-2780 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

## DATA



Western blot analysis of Ac-Histone H4 acetylation in untreated (A) and Scriptaid (sc-202807) treated (B) A549 whole cell lysates. Antibody tested include Ac-Histone H4 (Ser 1/Lys 5/Lys 8/Lys 12): sc-34263 (A,B). Note acetylation of Ac-Histone H4 in lane B.



Ac-Histone H4 (Ser 1/Lys 5/Lys 8/Lys 12): sc-34263. Immunofluorescence staining of methanol-fixed HeLa cells showing nuclear localization.

## SELECT PRODUCT CITATIONS

- Liao, W., et al. 2013. Effect of epigenetic histone modifications on E-cadherin splicing and expression in lung cancer. *Am. J. Cancer Res.* 3: 374-389.
- Ragu Varman, D. and Rajan, K.E. 2015. Environmental enrichment reduces anxiety by differentially activating serotonergic and neuropeptide Y (NPY)-ergic system in indian field mouse (*Mus booduga*): an animal model of post-traumatic stress disorder. *PLoS ONE* 10: e0127945.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.



Try **Ac-Histone H4 (E-5): sc-377520** or **Ac-Histone H4 (F-3): sc-377521**, our highly recommended monoclonal alternatives to Ac-Histone H4 (Ser 1/Lys 5/Lys 8/Lys 12).