# LTB4R2 (T-13): sc-34349



The Power to Question

### **BACKGROUND**

The P2Y receptor family consists of G protein-coupled receptors, which mediate the effects of extracellular nucleotides, primarily through the activation of phospholipase C. The P2Y receptors are important in the activation of leukocytes and platelets in response to inflammation or vascular damage. Leukotriene B4 receptor 1 (LTB4-R1), also designated P2Y purinoceptor 7, is a receptor for extracellular ATP, UTP and ADP. Through modulation of L-type calcium currents, LTB4-R1 is involved in the regulation of cardiac muscle contraction. It is also a receptor for leukotriene B4 (LTB4), a potent chemo-attractant involved in inflammation and immumne response. Leukotriene B4 receptor 2 (LTB4R2) is a ubiquitously expressed 358 amino acid receptor for LTB4. It is an integral membrane protein acting as a receptor for leukotrienes, is widely expressed and mediates chemotaxis of macrophages and granulocytes.

# **REFERENCES**

- Kamohara, M., et al. 2000. Molecular cloning and characterization of another leukotriene B4 receptor. J. Biol. Chem. 275: 27000-27004.
- Wang, S., et al. 2000. A novel hepatointestinal leukotriene B4 receptor. Cloning and functional characterization. J. Biol. Chem. 275: 40686-40694.
- Tryselius, Y., et al. 2000. Cloning and characterization of cDNA encoding a novel human leukotriene B4 receptor. Biochem. Biophys. Res. Commun. 274: 377-382.
- Nilsson, N.E., et al. 2000. Genomic organization of the leukotriene B4 receptor locus of human chromosome 14. Biochem. Biophys. Res. Commun. 274: 383-388.
- 5. Kato, K., et al. 2000. Cell-specific transcriptional regulation of human leukotriene B4 receptor gene. J. Exp. Med. 192: 413-420.
- Yokomizo, T., et al. 2000. A second leukotriene B4 receptor, BLT2. A new therapeutic target in inflammation and immunological disorders. J. Exp. Med. 192: 421-432.
- 7. Yokomizo, T., et al. 2001. Hydroxyeicosanoids bind to and activate the low affinity leukotriene B4 receptor, BLT2. J. Biol. Chem. 276: 12454-12459.

## **CHROMOSOMAL LOCATION**

Genetic locus: LTB4R2 (human) mapping to 14q11.2-q12; Ltb4r2 (mouse) mapping to 14 C1.

## **SOURCE**

LTB4R2 (T-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping within a cytoplasmic domain of LTB4R2 of human origin.

## **STORAGE**

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## **PRODUCT**

Each vial contains 200  $\mu g$  lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34349 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

# **APPLICATIONS**

LTB4R2 (T-13) is recommended for detection of LTB4R2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000); may cross-react with RIN1.

Suitable for use as control antibody for LTB4R2 siRNA (h): sc-45322, LTB4R2 siRNA (m): sc-45323, LTB4R2 shRNA Plasmid (h): sc-45322-SH, LTB4R2 shRNA Plasmid (m): sc-45323-SH, LTB4R2 shRNA (h) Lentiviral Particles: sc-45322-V and LTB4R2 shRNA (m) Lentiviral Particles: sc-45323-V.

Molecular Weight of LTB4R2: 42 kDa.

# **RECOMMENDED SECONDARY REAGENTS**

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat lgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat lgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat lgG-FITC: sc-2024 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### **PROTOCOLS**

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**