

## Sec15B (C-14): sc-34375

### BACKGROUND

Exocytosis, crucial in membrane trafficking, mediates hormone and neurotransmitter secretion out of the cell, as well as the incorporation of membrane proteins and lipids to the plasma membrane. It is crucial for cell-cell communication, cell growth and cell polarity. The exocyst complex is a multi-protein complex that consists of Sec3, Sec5, Sec6, Sec8, Sec10, Sec15, Exo70 and Exo84 and is essential for targeting exocytic vesicles to specific docking sites on the plasma membrane. The exocyst complex inhibits Tubulin polymerization *in vitro*, which implicates the exocyst in modulating microtubule dynamics underlying exocytosis. Sec15A (also designated Sec15L or Sec15p) and Sec15B (also designated Sec15L2) both belong to the Sec15 family of proteins. Sec15 can co-localize with Rab11 (a recycling endosome marker) and exhibits a GTP-dependent interaction with the Rab11 GTPase.

### REFERENCES

1. Wang, S., Liu, Y., Adamson, C.L., Valdez, G., Guo, W. and Hsu, S.C. 2004. The mammalian exocyst, a complex required for exocytosis, inhibits Tubulin polymerization. *J. Biol. Chem.* 279: 35958-35966.
2. Zhang, X.M., Ellis, S., Sriratana, A., Mitchell, C.A. and Rowe, T. 2004. Sec15 is an effector for the Rab11 GTPase in mammalian cells. *J. Biol. Chem.* 279: 43027-43034.
3. Hsu, S.C., TerBush, D., Abraham, M. and Guo, W. 2004. The exocyst complex in polarized exocytosis. *Int. Rev. Cytol.* 233: 243-265.
4. SWISS-PROT/TrEMBL (Q8TAG9). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

### CHROMOSOMAL LOCATION

Genetic locus: EXOC6B (human) mapping to 2p13.2; Exoc6b (mouse) mapping to 6 C3.

### SOURCE

Sec15B (C-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of Sec15B of human origin.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34375 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.

### APPLICATIONS

Sec15B (C-14) is recommended for detection of Sec15B of mouse and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Sec15B (C-14) is also recommended for detection of Sec15B in additional species, including equine, canine and porcine.

Suitable for use as control antibody for Sec15B siRNA (h): sc-45352, Sec15B siRNA (m): sc-45353, Sec15B shRNA Plasmid (h): sc-45352-SH, Sec15B shRNA Plasmid (m): sc-45353-SH, Sec15B shRNA (h) Lentiviral Particles: sc-45352-V and Sec15B shRNA (m) Lentiviral Particles: sc-45353-V.


Molecular Weight of Sec15B: 78 kDa.

Positive Controls: mouse brain extract: sc-2253.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### DATA



Sec15B (C-14): sc-34375. Western blot analysis of Sec15B expression in mouse brain tissue extract.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.