

## PAI-3 (M-17): sc-34495

### BACKGROUND

PAI-1, PAI-2 and PAI-3 (plasminogen activator inhibitor-1, -2 and -3) are members of the serpin serine proteinase inhibitor family. PAI-1 and PAI-2 regulate uPA (urokinase-type plasminogen activator) and TPA (tissue plasminogen activator), resulting in the inhibition of proteolytic activity. Members of the serpin family generally complex with their target proteinases, then disassociate slowly into cleaved species that fold into stable inactive forms. PAI-1 can fold into the inactive state without cleavage resulting in the latent form of PAI-1. Activity can be restored to the latent form of PAI-1 through denaturation and renaturation. PAI-2 occurs in secreted and cytosolic forms through facultative polypeptide translocation. PAI-3 inhibits plasminogen activators as well as activated protein C. PAI-3 is secreted in plasma, but is also expressed in liver.

### REFERENCES

- Riccio, A., Grimaldi, G., Verde, P., Sebastio, G., Boast, S. and Blasi, F. 1985. The human urokinase-plasminogen activator gene and its promoter. *Nucleic. Acids Res.* 13: 2759-2771.
- Belin, D., Wohlwend, A., Schleuning, W.D., Kruithof, E.K. and Vassalli, J.D. 1989. Facultative polypeptide translocation allows a single mRNA to encode the secreted and cytosolic forms of PLI2. *EMBO J.* 8: 3287-3294.
- Schmitt, M., Kanayama, N., Janicke, F., Hafter, R. and Graeff, H. 1991. Human tumor cell urokinase-type plasminogen activator (uPA): degradation of the proenzyme form (pro-uPA) by granulocyte elastase prevents subsequent activation by plasmin. *Adv. Exp. Med. Biol.* 297: 111-128.
- Mottonen, J., Strand, A., Symersky, J., Sweet, R.M., Danley, D.E., Geoghegan, K.F., Gerard, R.D. and Goldsmith, E.J. 1992. Structural basis of latency in PAI-1. *Nature* 355: 270-273.
- Niedbala, M.J. 1993. Cytokine regulation of endothelial cell extracellular proteolysis. *Agents Actions Suppl.* 42: 179-193.
- Schaefer, B.M., Stark, J.H., Fusenig, N.E., Todd, R.F. III and Kramer, M.D. 1995. Differential expression of urokinase-type plasminogen activator (uPA), its receptor (uPA-R), and inhibitor type-2 (PAI-2) during differentiation of keratinocytes in an organotypic coculture system. *Exp. Cell Res.* 220: 415-423.
- SWISS-PROT/TrEMBL (P05154). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

### CHROMOSOMAL LOCATION

Genetic locus: Serpina5 (mouse) mapping to 12 E.

### SOURCE

PAI-3 (M-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of PAI-3 of mouse origin.

### STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

### PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34495 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

### APPLICATIONS

PAI-3 (M-17) is recommended for detection of PAI-3 of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PAI-3 siRNA (m): sc-45417, PAI-3 shRNA Plasmid (m): sc-45417-SH and PAI-3 shRNA (m) Lentiviral Particles: sc-45417-V.

Molecular Weight of PAI-3: 46 kDa.

### RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.