

▶ LRF (P-14): sc-34511

BACKGROUND

LRF, formerly identified as Pokeman, is a poxvirus zinc finger (POZ) domain-containing transcription factor that influences cell differentiation. LRF (for leukemia/lymphoma related factor) is also known as zinc finger and BTB domain containing 7A, ZBTB7, TIP21, FBI1 and FBI-1. POZ-domain transcription factors contain a POZ or BTB type protein-protein interaction domain at their N-terminus and krüppel-type zinc fingers at their C-terminus. LRF is inducible during both murine and human preadipocyte differentiation and may contribute to adipogenesis through influencing the switch from cellular proliferation to terminal differentiation. LRF can associate with active chromatin and stimulate TAT-activated HIV-1 transcription through interactions with the HIV-1 long terminal repeat. 3T3L1 cells stably overexpressing LRF show a reduction in DNA synthesis and in expression of cyclin A, cyclin-dependent kinase 2 and p107.

CHROMOSOMAL LOCATION

Genetic locus: ZBTB7A (human) mapping to 19p13.3; Zbtb7a (mouse) mapping to 10 C1.

SOURCE

LRF (P-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping within an internal region of LRF of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available as TransCruz reagent for Gel Supershift and ChIP applications, sc-34511 X, 200 µg/0.1 ml.

Blocking peptide available for competition studies, sc-34511 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

LRF (P-14) is recommended for detection of LRF of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

LRF (P-14) is also recommended for detection of LRF in additional species, including canine, bovine and porcine.

Suitable for use as control antibody for LRF siRNA (h): sc-44574, LRF siRNA (m): sc-44575, LRF shRNA Plasmid (h): sc-44574-SH, LRF shRNA Plasmid (m): sc-44575-SH, LRF shRNA (h) Lentiviral Particles: sc-44574-V and LRF shRNA (m) Lentiviral Particles: sc-44575-V.

LRF (P-14) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of LRF: 72 kDa.

Positive Controls: Hs68 cell lysate: sc-2230.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



LRF (P-14): sc-34511. Immunofluorescence staining of formalin-fixed HepG2 cells showing nuclear localization.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **LRF (13E9): sc-33683** or **LRF (H-6): sc-393012**, our highly recommended monoclonal alternatives to LRF (P-14).