SANTA CRUZ BIOTECHNOLOGY, INC.

ABCA7 (P-16): sc-34559



BACKGROUND

ATP-binding cassette (ABC) transporters are an evolutionarily conserved family of widely-expressed proteins that use ATP hydrolysis to catalyze the transport of various molecules across extracellular and intracellular membranes. Eukaryotic ABC transporters are largely responsible for trafficking hydrophobic compounds either within the cell as part of a metabolic process, outside the cell for transport to other organs, or for secretion from the body. The cholesterolresponsive transporter, ABCA7, maps to human chromosome 19 and mouse chromosome 10 and has been reported as a candidate regulator of ceramide transport in epidermal lipid reorganization. High expression levels of ABCA7 have been reported in myelolymphatic tissues, reticuloendothelial cells, peripheral leukocytes, thymus, spleen and bone marrow. This expression pattern of the two alternatively-spliced isoforms also indicates an involvement in lipid homeostasis in cells of the immune system, though the complete role of ABCA7 is not yet known. Full-length type I ABCA7 has shown plasma membrane localization, while the type II splicing variant has shown expression predominantly in the endoplasmic reticulum.

REFERENCES

- Kaminski, W.E., et al. 2000. Genomic organization of the human cholesterolresponsive ABC transporter ABCA7: tandem linkage with the minor histocompatibility antigen HA-1 gene. Biochem. Biophys. Res. Commun. 278: 782-789.
- Broccardo, C., et al. 2001. Comparative analysis of the promoter structure and genomic organization of the human and mouse ABCA7 gene encoding a novel ABCA transporter. Cytogenet. Cell Genet. 92: 264-270.
- Ikeda, Y., et al. 2003. Post-transcriptional regulation of human ABCA7 and its function for the apoA-I-dependent lipid release. Biochem. Biophys. Res. Commun. 311: 313-318.
- Kielar, D., et al. 2003. Adenosine triphosphate binding cassette (ABC) transporters are expressed and regulated during terminal keratinocyte differentiation: a potential role for ABCA7 in epidermal lipid reorganization. J. Invest. Dermatol. 121: 465-474.

CHROMOSOMAL LOCATION

Genetic locus: ABCA7 (human) mapping to 19p13.3; Abca7 (mouse) mapping to 10 C1.

SOURCE

ABCA7 (P-16) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the N-terminus of ABCA7 of human origin.

PRODUCT

Each vial contains 200 μg lgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34559 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

ABCA7 (P-16) is recommended for detection of ABCA7, isoforms a and b of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for ABCA7 siRNA (h): sc-45431, ABCA7 siRNA (m): sc-45432, ABCA7 shRNA Plasmid (h): sc-45431-SH, ABCA7 shRNA Plasmid (m): sc-45432-SH, ABCA7 shRNA (h) Lentiviral Particles: sc-45431-V and ABCA7 shRNA (m) Lentiviral Particles: sc-45432-V.

Molecular Weight of ABCA7: 251 kDa.

Positive Controls: rat thymus extract: sc-2401.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.





ABCA7 (P-16): sc-34559. Western blot analysis of human recombinant ABCA7 fusion protein.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

MONOS Satisfation Guaranteed

Try ABCA7 (E-11): sc-377335 or ABCA7 (7A1-144): sc-130576, our highly recommended monoclonal alternatives to ABCA7 (P-16).