

# Dicer (I-15): sc-34601

## BACKGROUND

The mammalian Dicer is a type III RNase-related protein with orthologs in yeast, *Drosophila* and *Arabidopsis*. Dicer contains an RNA-helicase motif including a DEXH-box in its amino-terminus and an RNase motif in the carboxy-terminus. The gene encoding human Dicer maps to chromosome 14q32.13. Dicer is expressed in brain, heart, liver, lung, pancreas, kidney and placenta and functions in the RNA interference pathway. Dicer cleaves short hairpin RNA precursors of approximately 70 bp into 21-23 bp dsRNAs that selectively target the destruction of homologous RNAs. Dicer localizes to the cytoplasm of mammalian cells. Specifically, it co-localizes with calreticulin in the endoplasmic reticulum. Although the cleavage of RNA by Dicer is ATP-independent, the product release necessary for the rapid turnover of this enzyme may be attributed to ATP. Immunoprecipitation studies indicate Dicer forms a complex with PIWI domain of eIF2C translation initiation factors.

## REFERENCES

1. Matsuda, S., et al. 2000. Molecular cloning and characterization of a novel human gene (HERNA) which encodes a putative RNA-helicase. *Biochim. Biophys. Acta* 1490: 163-169.
2. Billy, E., et al. 2001. Specific interference with gene expression induced by long, double-stranded RNA in mouse embryonal teratocarcinoma cell lines. *Proc. Natl. Acad. Sci. USA* 98: 14428-14433.
3. Provost, P., et al. 2002. Ribonuclease activity and RNA binding of recombinant human Dicer. *EMBO J.* 21: 5864-5874.
4. Zhang, H., et al. 2002. Human Dicer preferentially cleaves dsRNAs at their termini without a requirement for ATP. *EMBO J.* 21: 5875-5885.
5. Paddison, P.J., et al. 2002. Short hairpin RNAs (shRNAs) induce sequence-specific silencing in mammalian cells. *Genes Dev.* 16: 948-958.
6. Doi, N., et al. 2004. Short-interfering RNA-mediated gene silencing in mammalian cells requires Dicer and eIF2C translation initiation factors. *Curr. Biol.* 13: 41-46.

## CHROMOSOMAL LOCATION

Genetic locus: DICER1 (human) mapping to 14q32.13; Dicer1 (mouse) mapping to 12 E.

## SOURCE

Dicer (I-15) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of Dicer of human origin.

## PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34601 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## APPLICATIONS

Dicer (I-15) is recommended for detection of Dicer of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Dicer (I-15) is also recommended for detection of Dicer in additional species, including equine, canine, porcine and avian.

Suitable for use as control antibody for Dicer siRNA (h): sc-40489, Dicer siRNA (m): sc-40490, Dicer shRNA Plasmid (h): sc-40489-SH, Dicer shRNA Plasmid (m): sc-40490-SH, Dicer shRNA (h) Lentiviral Particles: sc-40489-V and Dicer shRNA (m) Lentiviral Particles: sc-40490-V.

Molecular Weight of Dicer: 218 kDa.

Positive Controls: rat brain extract: sc-2392, Hep G2 cell lysate: sc-2227 or Y79 cell lysate: sc-2240.

## DATA



Dicer (I-15): sc-34601. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) or our catalog for detailed protocols and support products.


 MONOS  
Satisfaction  
Guaranteed

Try **Dicer (F-10): sc-136979** or **Dicer (E-7): sc-393328**, our highly recommended monoclonal alternatives to Dicer (I-15). Also, for AC, HRP, FITC, PE, Alexa Fluor® 488 and Alexa Fluor® 647 conjugates, see **Dicer (F-10): sc-136979**.