

TudorSN (C-17): sc-34753

BACKGROUND

TudorSN functions in the Pim-1 regulation of Myb activity and acts as a transcriptional activator of EBNA-2. TudorSN also interacts with EAV, NSP1, GTF2E1 and GTF2E2, and forms a ternary complex with Stat6 and POLR2A. The staphylococcal nuclease-like (SN)-domains directly interact with amino acids 1,099-1,758 of CBP. TudorSN plays an important role in the assembly of Stat6 transcriptome and stimulates IL-4-dependent transcription by mediating interaction between Stat6 and CBP.

CHROMOSOMAL LOCATION

Genetic locus: SND1 (human) mapping to 7q32.1; Snd1 (mouse) mapping to 6 A3.3.

SOURCE

TudorSN (C-17) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of TudorSN of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34753 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-34753 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

TudorSN (C-17) is recommended for detection of TudorSN of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TudorSN (C-17) is also recommended for detection of TudorSN in additional species, including equine, canine, bovine, porcine and avian.

Suitable for use as control antibody for TudorSN siRNA (h): sc-45514, TudorSN siRNA (m): sc-45515, TudorSN shRNA Plasmid (h): sc-45514-SH, TudorSN shRNA Plasmid (m): sc-45515-SH, TudorSN shRNA (h) Lentiviral Particles: sc-45514-V and TudorSN shRNA (m) Lentiviral Particles: sc-45515-V.

TudorSN (C-17) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TudorSN: 100 kDa.

Positive Controls: TudorSN (m): 293T Lysate: sc-124365, KNRK nuclear extract: sc-2141 or NIH/3T3 nuclear extract: sc-2138.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



TudorSN (C-17): sc-34753. Western blot analysis of TudorSN expression in Jurkat (A), KNRK (B) and NIH/3T3 (C) nuclear extracts and rat (D) and mouse (E) liver tissue extracts.



TudorSN (C-17): sc-34753. Western blot analysis of TudorSN expression in non-transfected: sc-117752 (A) and mouse TudorSN transfected: sc-124365 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Nalvarte, I., et al. 2010. Proteomics analysis of the estrogen receptor α receptosome. *Mol. Cell. Proteomics* 9: 1411-1422.
- Gao, X., et al. 2012. Tudor staphylococcal nuclease (Tudor-SN) participates in small ribonucleoprotein (snRNP) assembly via interacting with symmetrically dimethylated Sm proteins. *J. Biol. Chem.* 287: 18130-18141.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **TudorSN (F-5): sc-166676** or **TudorSN (C-9): sc-271590**, our highly recommended monoclonal alternatives to TudorSN (C-17).