

TudorSN (N-13): sc-34757

BACKGROUND

TudorSN functions in the Pim-1 regulation of Myb activity and acts as a transcriptional activator of EBNA-2. TudorSN also interacts with EAV, NSP1, GTF2E1 and GTF2E2, and forms a ternary complex with Stat6 and POLR2A. The staphylococcal nuclease-like (SN)-domains directly interact with amino acids 1099-1758 of CBP. TudorSN plays an important role in the assembly of Stat6 transcriptome and stimulates IL-4-dependent transcription by mediating interaction between Stat6 and CBP.

REFERENCES

1. Levenson, J.D., et al. 1998. Pim-1 kinase and p100 cooperate to enhance c-Myb activity. *Mol. Cell* 2: 417-425.
2. Tijms, M.A., et al. 2003. Equine arteritis virus non-structural protein 1, an essential factor for viral subgenomic mRNA synthesis, interacts with the cellular transcription J. *Gen. Virol.* 84: 2317-2322.
3. Pauku, K., et al. 2003. Tudor and nuclease-like domains containing protein p100 function as coactivators for signal transducer and activator of transcription 5. *Mol. Endocrinol.* 17: 1805-1814.
4. Broadhurst, M.K., et al. 2005. The p100 EBNA-2 coactivator: a highly conserved protein found in a range of exocrine and endocrine cells and tissues in cattle. *Biochim. Biophys. Acta* 1681: 126-133.
5. Valineva, T., et al. 2005. The transcriptional co-activator protein p100 recruits histone acetyltransferase activity to Stat6 and mediates interaction between the CREB-binding protein and Stat6. *J. Biol. Chem.* 280: 14989-14996.

CHROMOSOMAL LOCATION

Genetic locus: SND1 (human) mapping to 7q32.1.

SOURCE

TudorSN (N-13) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the N-terminus of TudorSN of human origin.

PRODUCT

Each vial contains 200 µg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34757 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

Available as TransCruz reagent for Gel Supershift and ChIP applications, sc-34757 X, 200 µg/0.1 ml.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

TudorSN (N-13) is recommended for detection of TudorSN of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

TudorSN (N-13) is also recommended for detection of TudorSN in additional species, including equine, canine and bovine.

Suitable for use as control antibody for TudorSN siRNA (h): sc-45514, TudorSN shRNA Plasmid (h): sc-45514-SH and TudorSN shRNA (h) Lentiviral Particles: sc-45514-V.

TudorSN (N-13) X TransCruz antibody is recommended for Gel Supershift and ChIP applications.


Molecular Weight of TudorSN: 100 kDa.

Positive Controls: Jurkat nuclear extract: sc-2132.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941. 3) Immunohistochemistry: use ImmunoCruz™: sc-2053 or ABC: sc-2023 goat IgG Staining Systems.

DATA



TudorSN (N-13): sc-34757. Immunoperoxidase staining of formalin fixed, paraffin-embedded human pancreas tissue showing cytoplasmic staining of Islets of Langerhans.

RESEARCH USE

For research use only, not for use in diagnostic procedures.