

I-TAC (E-14): sc-34785

BACKGROUND

IFN- γ -inducible T cell α chemoattractant (I-TAC), also designated β R1, H174, SCYB9B, Scyb11 (mouse), IP-9 or CXCL11, is a member of the CXC chemokine family and is expressed in IFN- γ -treated astrocytes, monocytes, keratinocytes, bronchial epithelial cells and neutrophils. The gene encoding I-TAC maps to human chromosome 4q21.2. I-TAC and two related proteins, IFN-induced protein of 10 kDa (IP-10) and monokine induced by IFN- γ (MIG), belong to the non-glutamate-leucine-arginine motif CXC chemokine family and act solely through the CXCR-3 receptor for potent attraction of T lymphocytes. I-TAC is assumed to be involved in inflammatory diseases characterized by the presence of activated T cells.

REFERENCES

- Mach, F., Sauty, A., Iarossi, A.S., Sukhova, G.K., Neote, K., Libby, P. and Luster, A.D. 1999. Differential expression of three T lymphocyte-activating CXC chemokines by human atheroma-associated cells. *J. Clin. Invest.* 104: 1041-1050.
- Tensen, C.P., Flier, J., Rampersad, S.S., Sampat-Sardjoepersad, S., Scheper, R.J., Boersma, D.M. and Willemze, R. 1999. Genomic organization, sequence and transcriptional regulation of the human CXCL11(1) gene. *Biochim. Biophys. Acta* 1446: 167-172.
- Meyer, M., Erdel, M., Duba, H.C., Werner, E.R. and Werner-Felmayer, G. 2000. Cloning, genomic sequence and chromosome mapping of Scyb11, the murine homologue of SCYB11 (alias β R1/H174/SCYB9B/I-TAC/IP-9/CXCL11). *Cytogenet. Cell Genet.* 88: 278-282.
- Mazanet, M.M., Neote, K. and Hughes, C.C. 2000. Expression of IFN- γ -inducible T cell α chemoattractant by human endothelial cells is cyclosporin A-resistant and promotes T cell adhesion: implications for cyclosporin A-resistant immune inflammation. *J. Immunol.* 164: 5383-5388.
- Albanesi, C., Scarponi, C., Sebastiani, S., Cavani, A., Federici, M., De Pita, O., Puddu, P. and Girolomoni, G. 2000. IL-4 enhances keratinocyte expression of CXCR-3 agonistic chemokines. *J. Immunol.* 165: 1395-1402.

CHROMOSOMAL LOCATION

Genetic locus: Cxcl11 (mouse) mapping to 5 E3.

SOURCE

I-TAC (E-14) is an affinity purified goat polyclonal antibody raised against a peptide mapping near the C-terminus of I-TAC of mouse origin.

PRODUCT

Each vial contains 200 μ g IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-34785 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

APPLICATIONS

I-TAC (E-14) is recommended for detection of I-TAC of mouse and rat origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for I-TAC siRNA (m): sc-39355, I-TAC shRNA Plasmid (m): sc-39355-SH and I-TAC shRNA (m) Lentiviral Particles: sc-39355-V.

Molecular Weight of I-TAC: 9 kDa.

RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

DATA



I-TAC (E-14): sc-34785. Western blot analysis of mouse recombinant I-TAC.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.



Try **I-TAC (R-15): sc-74094**, our highly recommended monoclonal alternative to I-TAC (E-14).