SANTA CRUZ BIOTECHNOLOGY, INC.

H-Ras (259): sc-35



BACKGROUND

The mammalian Ras (also designated v-Ha-Ras, Harvey rat sarcoma viral oncogene homolog, HRAS1, K-Ras, N-Ras, RASH1 or c-bas/has) gene family consists of the Harvey and Kirsten Ras genes (c-H-Ras1 and c-K-Ras2), an inactive pseudogene of each (c-H-Ras2 and c-K-Ras1) and the N-Ras gene. The three Ras oncogenes, H-Ras, K-Ras and N-Ras, encode proteins with GTP/ GDP binding and GTPase activity. Ras proteins alternate between an inactive form bound to GDP and an active form bound to GTP, activated by a guanine nucleotide-exchange factor (GEF) and inactivated by a GTPase-activating protein (GAP). Ras nomenclature originates from the characterization of human DNA sequences homologous to cloned DNA fragments containing oncogenic sequences of a type C mammalian retrovirus, the Harvey strain of murine sarcoma virus (HaMSV), derived from the rat. Under normal conditions, Ras family members influence cell growth and differentiation events in a subcellular membrane compartmentalization-based signaling system. Oncogenic Ras can deregulate processes that control both cell proliferation and apoptosis. The Ras superfamily of GTP hydrolysis-coupled signal transduction relay proteins can be subclassified into Ras, Rho, Rab and ARF families.

SOURCE

H-Ras (259) is a rat monoclonal antibody raised against full length Ras p21 of human origin.

PRODUCT

Each vial contains 200 μ g lgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin. Also available azide-free for blocking H-Ras binding of Raf, sc-35 L, 200 μ g/0.1 ml.

H-Ras (259) is available conjugated to agarose (sc-35 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-35 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-35 PE), fluorescein (sc-35 FITC), Alexa Fluor[®] 488 (sc-35 AF488), Alexa Fluor[®] 546 (sc-35 AF546), Alexa Fluor[®] 594 (sc-35 AF594) or Alexa Fluor[®] 647 (sc-35 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-35 AF680) or Alexa Fluor[®] 790 (sc-35 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-35 P, (100 μ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

H-Ras (259) is recommended for detection of antigenic determinants common to H-, K- and N-Ras p21 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), flow cytometry (1 μ g per 1 x 10⁶ cells) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Molecular Weight of H-Ras: 21 kDa.

Positive Controls: K-562 whole cell lysate: sc-2203, MCF7 whole cell lysate: sc-2206 or Jurkat whole cell lysate: sc-2204.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA





H-Ras (259): sc-35. Western blot analysis of H-Ras expression in K-562 (**A**), HeLa (**B**), Jurkat (**C**), NIH/3T3 (**D**), A-431 (**E**) and MCF7 (**F**) whole cell lysates.

H-Ras (259) AF488: sc-35 AF488. Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization.

SELECT PRODUCT CITATIONS

- Umanoff, H., et al. 1995. The murine N-Ras gene is not essential for growth and development. Proc. Natl. Acad. Sci. USA 92: 1709-1713.
- Zhou, H., et al. 2012. Ginkgolide B inhibits renal cyst development in *in vitro* and *in vivo* cyst models. Am. J. Physiol. Renal Physiol. 302: F1234-F1242.
- Basile, K.J., et al. 2013. *In vivo* MAPK reporting reveals the heterogeneity in tumoral selection of resistance to RAF inhibitors. Cancer Res. 73: 7101-7110.
- 4. Janardhan, S.V., et al. 2014. Primary murine CD4⁺ T cells fail to acquire the ability to produce effector cytokines when active Ras is present during Th1/Th2 differentiation. PLoS ONE 9: e112831.
- Vega, S.L., et al. 2015. Organizational metrics of interchromatin speckle factor domains: integrative classifier for stem cell adhesion & lineage signaling. Integr. Biol. 7: 435-446.
- 6. Saha, S.K., et al. 2016. KRT19 directly interacts with β -catenin/RAC1 complex to regulate NUMB-dependent NOTCH signaling pathway and breast cancer properties. Oncogene 36: 332-349.
- 7. García-Pérez, D., et al. 2017. Acute morphine, chronic morphine, and morphine withdrawal differently affect pleiotrophin, midkine, and receptor protein tyrosine phosphatase β/ζ regulation in the ventral tegmental area. Mol. Neurobiol. 54: 495-510.
- Tanaka, M., et al. 2018. The inhibitory NKR-P1B:Clr-b recognition axis facilitates detection of oncogenic transformation and cancer immunosurveillance. Cancer Res. 78: 3589-3603.
- Wu, J., et al. 2019. Novel compound cedrelone inhibits hepatocellular carcinoma progression via PBLD and Ras/Rap1. Exp. Ther. Med. 18: 4209-4220.

RESEARCH USE

For research use only, not for use in diagnostic procedures.