

GSK-3 β siRNA (m): sc-35525

BACKGROUND

Glycogen synthase kinase 3, or GSK-3, is a serine/threonine, proline-directed kinase involved in a diverse array of signaling pathways, including glycogen synthesis and cellular adhesion, and has been implicated in Alzheimer's disease. Two forms of GSK-3, designated GSK-3 α and GSK-3 β , have been identified and differ in their subcellular localization. Tau, a microtubule-binding protein which serves to stabilize microtubules in growing axons, is found to be hyper-phosphorylated in paired helical filaments (PHF), the major fibrous component of neurofibrillary lesions associated with Alzheimer's disease. Hyperphosphorylation of Tau is thought to be the critical event leading to the assembly of PHF. Six Tau protein isoforms have been identified, all of which are phosphorylated by GSK-3. This presents the possibility that miscues in GSK-3 signaling contribute to the onset of Alzheimer's disease.

CHROMOSOMAL LOCATION

Genetic locus: Gsk3b (mouse) mapping to 16 B3.

PRODUCT

GSK-3 β siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GSK-3 β shRNA Plasmid (m): sc-35525-SH and GSK-3 β shRNA (m) Lentiviral Particles: sc-35525-V as alternate gene silencing products.

For independent verification of GSK-3 β (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-35525A, sc-35525B and sc-35525C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GSK-3 β siRNA (m) is recommended for the inhibition of GSK-3 β expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

GSK-3 β (E-11): sc-377213 is recommended as a control antibody for monitoring of GSK-3 β gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GSK-3 β gene expression knockdown using RT-PCR Primer: GSK-3 β (m)-PR: sc-35525-PR (20 μ l, 489 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

SELECT PRODUCT CITATIONS

- Zhou, F., et al. 2008. The association of GSK-3 β with E2F1 facilitates nerve growth factor-induced neural cell differentiation. *J. Biol. Chem.* 283: 14506-14515.
- He, P. and Shen, Y. 2009. Interruption of β -catenin signaling reduces neurogenesis in Alzheimer's disease. *J. Neurosci.* 29: 6545-6557.
- Zou, C., et al. 2011. LPS impairs phospholipid synthesis by triggering β -transducin repeat-containing protein (β -TrCP)-mediated polyubiquitination and degradation of the surfactant enzyme acyl-CoA:lysophosphatidylcholine acyltransferase 1 (LPCAT1). *J. Biol. Chem.* 286: 2719-2727.
- Kim, K.J., et al. 2012. A chemical genomics screen to discover genes that modulate neural stem cell differentiation. *J. Biomol. Screen.* 17: 129-139.
- Lu, J., et al. 2013. Formaldehyde induces hyperphosphorylation and polymerization of Tau protein both *in vitro* and *in vivo*. *Biochim. Biophys. Acta* 1830: 4102-4116.
- Kim, S., et al. 2015. Endoplasmic reticulum stress-induced IRE1 α activation mediates cross-talk of GSK-3 β and XBP-1 to regulate inflammatory cytokine production. *J. Immunol.* 194: 4498-4506.
- L'Episcopo, F., et al. 2016. GSK-3 β -induced Tau pathology drives hippocampal neuronal cell death in Huntington's disease: involvement of astrocyte-neuron interactions. *Cell Death Dis.* 7: e2206.
- Yushan, R., et al. 2018. Isoliquiritigenin inhibits mouse S180 tumors with a new mechanism that regulates autophagy by GSK-3 β /TNF- α pathway. *Eur. J. Pharmacol.* 838: 11-22.
- Breit, A., et al. 2018. Insulin-like growth factor-1 acts as a zeitgeber on hypothalamic circadian clock gene expression via glycogen synthase kinase-3 β signalling. *J. Biol. Chem.* 293: 17278-17290.
- Lee, H.J., et al. 2020. TLR7 stimulation with imiquimod induces selective autophagy and controls *Mycobacterium tuberculosis* growth in mouse macrophages. *Front. Microbiol.* 11: 1684.

RESEARCH USE

For research use only, not for use in diagnostic procedures.