

West Nile - Innovator® (FORT DODGE ANIMAL HEALTH)

Introduction **Label**

Wyeth is a wholly owned subsidiary of Pfizer Inc. - to learn more please visit www.pfizer.com
Fort Dodge is now part of Pfizer Animal Health - to learn more please visit www.pfizeranimalhealth.com

FORT DODGE ANIMAL HEALTH

Division of Wyeth

800-5TH STREET N.W., P.O. BOX 518, FORT DODGE, IA, 50501

Telephone: 515-955-4600
Fax: 515-302-2693
Order Desk Telephone: 800-685-5656
Order Desk Fax: 800-846-8626
Professional Services: 800-533-8536
Website: www.wyeth.com/divisions/fort_dodge.asp
Order Desk Email: fdorder@FDAH.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

WEST NILE - INNOVATOR®

Fort Dodge

West Nile Virus Vaccine

Killed Virus

INDICATIONS: For intramuscular vaccination of healthy horses 10 months of age or older as an aid in the prevention of viremia caused by West Nile Virus. Efficacy was demonstrated in horses that received two doses of the vaccine and were challenged one year post-vaccination with West Nile Virus. The vaccine protected 95% of the vaccinates against viremia, while 82% of the controls were viremic.

Owners should be advised that vaccinated horses may develop antibody to West Nile Virus, which may affect their ability to export the animal.

DOSE: Inject one 1 mL dose intramuscularly using aseptic technique. Administer a second 1 mL dose 3 to 6 weeks after the first dose. Revaccinate annually with one 1 mL dose. Early revaccination may be advisable when horses are faced with an outbreak or with other conditions which might make heavy exposure likely.

CAUTION: Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. Use entire contents when first opened. In some instances, transient local reactions may occur at the injection site. Do not vaccinate within 21 days before slaughter. In case of anaphylactoid reaction, administer epinephrine.

Thimerosal, neomycin and polymyxin B added as preservatives.

For Veterinary Use Only

U.S. Pat. Nos. 7,153,513; 7,445,787 and patents pending

© 2000 Fort Dodge Animal Health. All Rights Reserved.

Fort Dodge Animal Health, A division of Wyeth, Fort Dodge, Iowa 50501 USA

U.S. Vet. License No. 112

12 x 1 mL syringes		16661J
10 x 10 mL	100 Doses	16666E

NAC No.: 10032720