

Rumatel® Pellets

Durvet

Type C Medicated Feed

Indications:

Goats: For the removal and control of mature gastrointestinal nematode infections of goats including *Haemonchus contortus*, *Ostertagia (Teladorsagia) circumcincta*, and *Trichostrongylus axei*.
Cattle: For the removal and control of mature gastrointestinal nematode infections including stomach worms (*Haemonchus spp.*, *Ostertagia spp.*, *Trichostrongylus spp.*), worms of the small intestine (*Cooperia spp.*, *Trichostrongylus spp.*, *Nematodirus spp.*), and worms of the large intestine (*Oesophagostomum radiatum*).

Active Drug Ingredient:

Morantel Tartrate 4.45 g/lb.

Guaranteed Analysis:

Crude Protein not less than 15.0%
Crude Fat not less than 3.0%
Crude Fiber not more than 12.0%

Ingredients:

Wheat Middlings, Dehydrated Alfalfa Meal, Calcium Carbonate, Extruded Whole Soybeans, Calcium Propionic and Lignin Sulfonate.

Directions For Use:

Use a single therapeutic treatment in goats and cattle by feeding 0.1 pounds (.44 g morantel tartrate) of medicated ration per 100 pounds of body weight. May be fed as the sole ration, or mixed with 1 to 2 parts of complete feed or as a top dress. When used as a top dress, the medication as well as the underlying feed should be evenly distributed. Withhold feed overnight prior to treatment to insure the ration will be readily consumed. Group by size for optimum efficacy. Fresh water should be available at all times. When medicated feed is all consumed, resume normal feeding. Conditions of constant worm exposure may require retreatment within 2 to 4 weeks. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasites.

Caution:

Goats: Consult veterinarian before using in severely debilitated animals. Do not mix in feeds containing Bentonite.

Cattle: Consult veterinarian before using in severely debilitated animals. Do not mix in feeds containing Bentonite.

Warning:

Goats: Do not treat goats within 30 days of slaughter. No milk discard required following use in dairy goats.

Cattle: Do not treat cattle within 14 days of slaughter. No milk discard required following use in dairy cattle.