

Tylan® 200 Injection (ELANCO ANIMAL HEALTH)

ELANCO ANIMAL HEALTH

A Division of Eli Lilly & Co.

2500 INNOVATION WAY, GREENFIELD, IN, 46140

Main Switchboard:
Customer Service:
Small Animal Product Customer Service:
Technical Services:
Fax:
Website:
Email:

317-433-4800
317-276-1262
888-545-5973
800-428-4441
317-276-2270
www.elanco.com
elanco@elanco.com



Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the USA product label or package insert.

TYLAN® 200 INJECTION

Elanco

Tylosin Injection

200 mg per mL

For Use in Cattle and Swine Only

An Antibiotic

500 mL - Use automatic syringe equipment only

Indications: In Beef Cattle and Non-lactating Dairy Cattle, Tylan 200 Injection is indicated for use in the treatment of bovine respiratory complex (shipping fever, pneumonia) usually associated with *Pasteurella multocida* and *Actinomyces pyogenes*; foot rot (necrotic pododermatitis) and calf diphtheria caused by *Fusobacterium necrophorum* and metritis caused by *Actinomyces pyogenes*.

In Swine, Tylan 200 Injection is indicated for use in the treatment of swine arthritis caused by *Mycoplasma hyosynoviae*; swine pneumonia caused by *Pasteurella spp.*; swine erysipelas caused by *Erysipelothrix rhusiopathiae*; swine dysentery associated with *Treponema hyodysenteriae* when followed by appropriate medication in the drinking water and/or feed.

Each mL contains 200 mg of tylosin activity (as tylosin base) in 50 percent propylene glycol with 4 percent benzyl alcohol and water for injection.

ADMINISTRATION AND DOSAGE: Tylan 200 Injection is administered intramuscularly.

BEEF CATTLE AND NON-LACTATING DAIRY CATTLE-Inject intramuscularly 8 mg per pound of body weight one time daily (1 mL per 25 pounds). Treatment should be continued for 24 hours following remission of disease signs, not to exceed 5 days. Do not inject more than 10 mL per site.

SWINE-Inject intramuscularly 4 mg per pound of body weight (1 mL per 50 pounds) twice daily. Treatment should be continued for 24 hours following remission of disease signs, not to exceed 3 days. Do not inject more than 5 mL per site.

Read accompanying directions fully before use.

CAUTION: Do not mix Tylan 200 Injection with other injectable solutions as this may cause a precipitation of the active ingredients. Do not administer to horses or other equines. Injection of tylosin in equines has been fatal.

PRECAUTION: Adverse reactions, including shock and death may result from overdosage in baby pigs. Do not attempt injection into pigs weighing less than 25 pounds (0.5 mL) with the common syringe. It is recommended that Tylan 50 Injection be used in pigs weighing less than 25 pounds.

If tylosin medicated drinking water is used as a follow-up treatment for swine dysentery the animal should thereafter receive feed containing 40 to 100 g of tylosin per ton for 2 weeks to assure depletion of tissue residues.

NOT FOR HUMAN USE

WARNING: Discontinue use in cattle 21 days before slaughter. Discontinue use in swine 14 days before slaughter. Do not use in lactating dairy cattle.

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Store at 72°F (22°C) or below.

*Elanco®, Tylan® and the diagonal color bar are trademarks of Eli Lilly and Company.

Restricted Drug (California) - Use Only as Directed.

NADA 12-965, Approved by FDA

To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

Manufactured for: Elanco Animal Health, A Division of Eli Lilly and Company, Indianapolis, IN 46285, U.S.A.

250 mL	AH 0206-38W	YL9510DEAMB (V01-08-2009)
500 mL Professional Size Use automatic syringe equipment only	AH 0206-25X	YL9520DEAMB (V01-08-2009)

NAC No.: 10310134