

Hu-CD19/CD69/CD45

3 Color FCM Reagent: sc-3607



BACKGROUND

Human CD19/CD69/CD45: sc-3607 is a direct immunofluorescence reagent formatted to identify activated B lymphocytes (1) in peripheral blood. CD19 is present on human B lymphocytes during all stages of B cell maturation, but is lost on plasma cells (2). In normal peripheral blood, CD69 is variably expressed on lymphocytes (3). Upon activation, CD69 expression increases on T, B, and NK lymphocytes (4). CD45 is a major leukocyte cell surface molecule (5). The total population of T lymphocytes and B lymphocytes are used to characterize and monitor some forms of immunodeficiency and autoimmune disease (6,7).

Antigen Expression	Cell Type Identified
CD3- CD19+	Total B Lymphocytes
CD3- CD19+ CD69+	Activated B Lymphocytes

STORAGE

Store at 4° C. Do not freeze. Stable for one year from the date of shipment. Protect reagents from prolonged exposure to light.

PRODUCT

Supplied in 1.0 ml of PBS containing 0.1% azide and 0.1% gelatin. Sufficient for 50 tests. This product has been titrated for optimal performance. Recommended use is 20 uL per test (1x10⁶ cells). **For research use only. Not for use in diagnostic procedures.**

INSTRUMENT

Human CD19/CD69/CD45: sc-3607 is recommended for use with either a single or dual laser Flow Cytometer fitted with appropriate acquisition and analysis software, such as the FACSCalibur™ Flow Cytometer fitted with CellQuest™ Software by Becton Dickinson.

The flow cytometer must be equipped with a 488 nm laser and must be capable of detecting light scatter (forward and side) and three-color fluorescence with emission detectable in three ranges: 515-545 nm, 562-607 nm and >650 nm, and it must be able to threshold and discriminate using the >650 channel.

Antigen	Clone	Isotype	Label*	Detection Range (nm)
CD19	SJ25C1	IgG ₁	FITC	515-545
CD69	FN50	IgG ₁	PE	562-607
CD45	2D-1	IgG ₁	PE-Cy5	>650

*Fluorescent labels include FITC: Fluorescein isothiocyanate; PE: phycoerythrin; PE-Cy5: phycoerythrin-cyanin 5.

ISOTYPE CONTROL

sc-3607 CON (IgG₁ FITC/IgG₁ PE/IgG₁ PE-Cy5) is the isotype matched negative control for this system and is suitable for 50 tests.

REFERENCES

- Testi, R., D'Ambrosio, D., De Maria, R., and Santoni, A. 1994. The CD69 receptor: a multipurpose cell-surface trigger for hematopoietic cells. *Immunol. Today* **15**: 479-483.
- Dörken, B., Möller, P., Pezzutto, A., Schwartz-Albiez, R., and Moldenhauer, G. B-cell antigens: CD19. In: Knapp, W., Dörken, B., Gilks, W.R. *et al.* eds. *Leucocyte Typing IV: White Cell Differentiation Antigens*. New York, NY: Oxford University Press; 1989: 34-36.
- Schwartz, R., Biedobitek, G., and Stein, H. Cluster report: CD69. Knapp, W., Dörken, B., Gilks, W.R., *et al.* eds. *Leucocyte Typing IV: White Cell Differentiation Antigens*. New York, NY: Oxford University Press; 1989: 428-432.
- Testi, R., Phillips, J.H., and Lanier, L.L. 1988. Constitutive expression of a phosphorylated activation antigen (Leu 23) by CD3bright human thymocytes. *J. Immunol.* **141**: 2557-2563.
- Charbonneau, H., Tonks, N.K., Walsh, K.A., and Fischer, E.H. 1988. The leukocyte common antigen (CD45): a putative receptor-linked protein tyrosine phosphatase. *Proc. Natl. Acad. Sci. USA* **85**: 7182-7186.
- Foucar, K. and Goeken, J.A. 1982. Clinical Applications of immunologic techniques to the diagnosis of lymphoproliferative and immunodeficiency disorders. *Lab. Med.* **13**: 403-413.
- Smolen, J.S., Chused, T.M., Leiserson, W.M., Reeves, J.P., Alling, D., and Steinberg, A.D. 1982. Heterogeneity of immunoregulatory T-cell subsets in systemic lupus erythematosus. Correlation with clinical features. *Am. J. Med.* **72**: 783-790.