

Mo-CD8/CD38/CD3

3 Color FCM Reagent: sc-3615



BACKGROUND

Mouse CD8/CD38/CD3: sc-3615 is a direct immunofluorescence reagent formatted to identify and determine the percentage of activated mature T lymphocytes and suppressor/cytotoxic T lymphocytes in erythrocyte-lysed whole blood, based on cell-surface antigen expression. CD3 identifies T lymphocytes and non-covalently associates with either α/β or γ/δ TCR (1). CD8 identifies suppressor/cytotoxic T lymphocytes and binds class I MHC molecules, which enhances the activation of resting T lymphocytes (2). CD3+CD8+ and CD3+CD4+ percentages are used to characterize and monitor some forms of immunodeficiency and autoimmune disease (3,4). CD38 is expressed on activated T and NK lymphocytes (5) and may play a role in regulating humoral immune responses (6).

Antigen Expression	Cell Type Identified
CD3+	Mature T Cells
CD3+ CD8+	Suppressor/Cytotoxic T Cells
CD3+ CD8+ CD38+	Activated S/C T Cells

STORAGE

Store at 4° C. Do not freeze. Stable for one year from the date of shipment. Protect reagents from prolonged exposure to light.

PRODUCT

Supplied in 1.0 ml of PBS containing 0.1% azide and 0.1% gelatin. Sufficient for 50 tests. This product has been titrated for optimal performance. Recommended use is 20 μ L per test (1x10⁶ cells). **For research use only. Not for use in diagnostic procedures.**

INSTRUMENT

Mouse CD8/CD38/CD3: sc-3615 is recommended for use with either a single or dual laser Flow Cytometer fitted with appropriate acquisition and analysis software, such as the FACSCalibur™ Flow Cytometer fitted with CellQuest™ Software by Becton Dickinson.

The flow cytometer must be equipped with a 488 nm laser and must be capable of detecting light scatter (forward and side) and three-color fluorescence with emission detectable in three ranges: 515-545 nm, 562-607 nm and >650 nm, and it must be able to threshold and discriminate using the >650 channel.

Antigen	Clone	Isotype	Label*	Detection Range (nm)
CD8	53-6.7	rat IgG _{2a}	FITC	515-545
CD38	90	rat IgG _{2a}	PE	562-607
CD3	145-2C11	Armenian Hamster IgG	PE-Cy5	>650

*Fluorescent labels include FITC: Fluorescein isothiocyanate; PE: phycoerythrin; PE-Cy5: phycoerythrin-cyanin 5.

ISOTYPE CONTROL

sc-3615 CON (rat IgG_{2a} FITC/rat IgG_{2a} PE/Armenian Hamster IgG PE-Cy5) is the isotype matched negative control for this system and is suitable for 50 tests.

REFERENCES

- Exley, M., Terhorst, C., and Wileman, T. 1991. Structure, assembly and intracellular transport of the T cell receptor for antigen. *Semin. Immunol.* 3: 283-297.
- Gallagher, P.F., Fazekas de St. Groth, B., and Miller, J.F. 1989. CD4 and CD8 molecules can physically associate with the same T-cell receptor. *Proc. Natl. Acad. Sci. USA* 86: 10044-10048.
- Foucar, K. and Goeken, J.A. 1982. Clinical Applications of immunologic techniques to the diagnosis of lymphoproliferative and immunodeficiency disorders. *Lab. Med.* 13: 403-413.
- Smolen, J.S., Chused, T.M., Leiserson, W.M., Reeves, J.P., Alling, D., and Steinberg, A.D. 1982. Heterogeneity of immunoregulatory T-cell subsets in systemic lupus erythematosus. Correlation with clinical features. *Am. J. Med.* 2: 783-790.
- Sconocchia, G., Titus, J.A., Mazzoni, A., Visintin, A., Pericle, F., Hicks, S.W., Malavasi, F., and Segal, D.M. 1999. CD38 triggers cytotoxic responses in activated human natural killer cells. *Blood* 94: 3864-3871.
- Cockayne, D.A., Muchamuel, T., Grimaldi, J.C., Muller-Steffner, H., Randall, T.D., Lund, F.E., Murray, R., Schuber, F., and Howard, M.C. 1998. Mice deficient for the ecto-nicotinamide adenine dinucleotide glycohydrolase CD38 exhibit altered humoral immune responses. *Blood* 92: 1324-1333.