

Rac 1 siRNA (h): sc-36351

BACKGROUND

A large number of low molecular weight, GTP binding proteins of the Ras superfamily have been identified. These proteins regulate many fundamental processes in all eukaryotic cells such as growth, vesicle traffic and cytoskeletal organization. GTPase-activating proteins (GAPs) accelerate the intrinsic rate of GTP hydrolysis of Ras-related proteins, resulting in downregulation of their active form. Two proteins in this family, Rac 1 and Rac 2, are 92% identical and share GTP binding and GTP hydrolysis motifs with other members of the Ras superfamily. Rac 1 is expressed in a large number of different cell types. Rac 2 is primarily expressed only in myeloid cells and has been reported to be a regulatory component of the human neutrophil NADPH oxidase.

CHROMOSOMAL LOCATION

Genetic locus: RAC1 (human) mapping to 7p22.1.

PRODUCT

Rac 1 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rac 1 shRNA Plasmid (h): sc-36351-SH and Rac 1 shRNA (h) Lentiviral Particles: sc-36351-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rac 1 siRNA (h) is recommended for the inhibition of Rac 1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

GENE EXPRESSION MONITORING

Rac 1/2/3 (G-2): sc-514583 is recommended as a control antibody for monitoring of Rac 1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rac 1 gene expression knockdown using RT-PCR Primer: Rac 1 (h)-PR: sc-36351-PR (20 μ l, 535 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Li, Q., et al. 2005. Interleukin-1 β induction of NF κ B is partially regulated by H₂O₂-mediated activation of NF κ B-inducing kinase. *J. Biol. Chem.* 281: 1495-1505.
- Pielage, J.F., et al. 2008. RNAi screen reveals an Abl kinase-dependent host cell pathway involved in *Pseudomonas aeruginosa* internalization. *PLoS Pathog.* 4: e1000031.
- Lyons, L.S., et al. 2008. Ligand-independent activation of androgen receptors by Rho GTPase signaling in prostate cancer. *Mol. Endocrinol.* 22: 597-608.
- Usatyuk, P.V., et al. 2009. Phospholipase D-mediated activation of IQGAP1 through Rac 1 regulates hyperoxia-induced p47-phox translocation and reactive oxygen species generation in lung endothelial cells. *J. Biol. Chem.* 284: 15339-15352.
- Liu, Z., et al. 2009. The Rho-specific guanine nucleotide exchange factor Dbs regulates breast cancer cell migration. *J. Biol. Chem.* 284: 15771-15780.
- Meissner, M., et al. 2009. Inhibition of Rac 1 GTPase downregulates vascular endothelial growth factor receptor-2 expression by suppressing Sp1-dependent DNA binding in human endothelial cells. *Exp. Dermatol.* 18: 863-869.
- Seidel, P., et al. 2010. DMF inhibits PDGF-BB induced airway smooth muscle cell proliferation through induction of Heme Oxygenase 1. *Respir. Res.* 11: 145.
- Visvikis, O., et al. 2011. *Escherichia coli* producing CNF1 toxin hijacks Tollip to trigger Rac 1-dependent cell invasion. *Traffic* 12: 579-590.
- Tugizov, S.M., et al. 2013. Epstein-Barr virus transcytosis through polarized oral epithelial cells. *J. Virol.* 87: 8179-8194.
- Han, R., et al. 2019. C3a and suPAR drive versican V1 expression in tubular cells of focal segmental glomerulosclerosis. *JCI Insight* 4: e122912.
- Zhang, L., et al. 2020. Rab 23 promotes hepatocellular carcinoma cell migration via Rac 1/TGF- β signaling. *Pathol. Oncol. Res.* 26: 301-306.

RESEARCH USE

For research use only, not for use in diagnostic procedures.