

DDO (H-6): sc-365135

BACKGROUND

DDO (D-aspartate oxidase), also known as DASOX, is a 341 amino acid protein that localizes to peroxisomes and exists as two alternatively spliced isoforms, designated DDO-1 and DDO-2. Using FAD or 6-hydroxyflavin adenine dinucleotide as cofactors, DDO functions as a peroxisomal flavoprotein that selectively catalyzes the oxidative deamination of D-aspartate and N-methyl D-aspartate. Human DDO shares 86% sequence similarity with its bovine counterpart, suggesting a conserved role between species. The gene encoding DDO maps to human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

REFERENCES

1. Barker, R.F. and Hopkinson, D.A. 1977. The genetic and biochemical properties of the D-amino acid oxidases in human tissues. *Ann. Hum. Genet.* 41: 27-42.
2. Van Veldhoven, P.P., et al. 1991. D-aspartate oxidase, a peroxisomal enzyme in liver of rat and man. *Biochim. Biophys. Acta* 1073: 203-208.
3. Nagasaki, H. 1994. Gender-related differences of mouse liver D-aspartate oxidase in the activity and response to administration of D-aspartate and peroxisome proliferators. *Int. J. Biochem.* 26: 415-423.
4. Simonic, T., et al. 1997. cDNA cloning and expression of the flavoprotein D-aspartate oxidase from bovine kidney cortex. *Biochem. J.* 322: 729-735.
5. Setoyama, C. and Miura, R. 1997. Structural and functional characterization of the human brain D-aspartate oxidase. *J. Biochem.* 121: 798-803.

CHROMOSOMAL LOCATION

Genetic locus: DDO (human) mapping to 6q21; Ddo (mouse) mapping to 10 B1.

SOURCE

DDO (H-6) is a mouse monoclonal antibody raised against amino acids 151-320 mapping near the C-terminus of DDO of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

DDO (H-6) is available conjugated to agarose (sc-365135 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-365135 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-365135 PE), fluorescein (sc-365135 FITC), Alexa Fluor® 488 (sc-365135 AF488), Alexa Fluor® 546 (sc-365135 AF546), Alexa Fluor® 594 (sc-365135 AF594) or Alexa Fluor® 647 (sc-365135 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-365135 AF680) or Alexa Fluor® 790 (sc-365135 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

DDO (H-6) is recommended for detection of DDO of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for DDO siRNA (h): sc-77101, DDO siRNA (m): sc-77102, DDO shRNA Plasmid (h): sc-77101-SH, DDO shRNA Plasmid (m): sc-77102-SH, DDO shRNA (h) Lentiviral Particles: sc-77101-V and DDO shRNA (m) Lentiviral Particles: sc-77102-V.


Molecular Weight of DDO: 37 kDa.

Positive Controls: DDO (m): 293T Lysate: sc-119700, Hep G2 cell lysate: sc-2227 or rat liver extract: sc-2395.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



DDO (H-6): sc-365135. Western blot analysis of DDO expression in non-transfected: sc-117752 (A) and mouse DDO transfected: sc-119700 (B) 293T whole cell lysates.



DDO (H-6): sc-365135. Western blot analysis of DDO expression in Hep G2 whole cell lysate (A) and rat liver tissue extract (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.