

creatine kinase-M (H-9): sc-365141

BACKGROUND

Creatine kinases (CKs) are a large family of isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. Creatine kinases provide the energy of phosphate hydrolysis necessary to drive the normal function of many cellular systems including muscle, electrocytes, retina photoreceptor cells, brain cells, kidney, salt glands, myometrium, placenta, pancreas, thymus, thyroid, intestinal epithelial cells, endothelial cells, cartilage and bone cells, macrophages, blood platelets, and tumor and cancer cells. Human cytoplasmic creatine kinase-B, also designated CK-B and BCK, is a 381 amino acid, brain tissue-specific isoform of creatine kinase. Human cytoplasmic creatine kinase-M (CK-M, MCK) is a muscle tissue-specific isoform of creatine kinase. Human cytoplasmic creatine kinase-Mi (Mi-CK, MtCK) is a 416 amino acid mitochondrial-specific isoform of creatine kinase. Cytosolic creatine kinases are important in the energetic regulation of Ca²⁺-pumps and in the maintenance of Ca²⁺-homeostasis.

REFERENCES

1. Mariman, E.C., et al. 1987. Structure and expression of the human creatine kinase B gene. *Genomics* 1: 126-137.
2. Nigro, J.M., et al. 1987. cDNA cloning and mapping of the human creatine kinase M gene to 19q13. *Am. J. Hum. Genet.* 40: 115-125.
3. Haas, R.C., et al. 1989. Isolation and characterization of the gene and cDNA encoding human mitochondrial creatine kinase. *J. Biol. Chem.* 264: 2890-2897.
4. Mariman, E.C., et al. 1989. Complete nucleotide sequence of the human creatine kinase B gene. *Nucleic Acids Res.* 17: 6385.
5. Wallimann, T., et al. 1994. Creatine kinase in non-muscle tissues and cells. *Mol. Cell. Biochem.* 133-134: 193-220.
6. Wallimann, T., et al. 1998. Some new aspects of creatine kinase (CK): compartmentation, structure, function and regulation for cellular and mitochondrial bioenergetics and physiology. *Biofactors* 8: 229-234.

CHROMOSOMAL LOCATION

Genetic locus: CKM (human) mapping to 19q13.32; Ckm (mouse) mapping to 7 A3.

SOURCE

creatine kinase-M (H-9) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 9-27 near the N-terminus of creatine kinase-M of human origin.

PRODUCT

Each vial contains 200 µg IgG₃ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365141 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

creatine kinase-M (H-9) is recommended for detection of creatine kinase-M chain of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


creatine kinase-M (H-9) is also recommended for detection of creatine kinase-M chain in additional species, including equine.

Suitable for use as control antibody for creatine kinase-M siRNA (h): sc-35109, creatine kinase-M siRNA (m): sc-35110, creatine kinase-M siRNA (r): sc-270230, creatine kinase-M shRNA Plasmid (h): sc-35109-SH, creatine kinase-M shRNA Plasmid (m): sc-35110-SH, creatine kinase-M shRNA Plasmid (r): sc-270230-SH, creatine kinase-M shRNA (h) Lentiviral Particles: sc-35109-V, creatine kinase-M shRNA (m) Lentiviral Particles: sc-35110-V and creatine kinase-M shRNA (r) Lentiviral Particles: sc-270230-V.


Molecular Weight of creatine kinase-M: 43 kDa.

Positive Controls: human skeletal muscle extract: sc-363776, human heart extract: sc-363763 or A-673 cell lysate: sc-2414.

DATA



creatine kinase-M (H-9): sc-365141. Western blot analysis of creatine kinase-M expression in human skeletal muscle (A) and human heart (B) tissue extracts.



creatine kinase-M (H-9): sc-365141. Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes (A). Immunofluorescence staining of methanol-fixed HeLa cells showing cytoplasmic localization (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.