

3pK (C-4): sc-365148

BACKGROUND

The MAPKAP kinases (for MAP kinase activated protein kinases) are a group of MAP kinase substrates which are themselves kinases. In response to activation, the MAP kinases phosphorylate downstream components on a consensus Pro-X-Ser/Thr-Pro motif. Several kinases that contain this motif have been identified and serve as substrates for the ERK and p38 MAP kinases. These include the serine/threonine kinases Rsk-1 (also designated MAPKAP kinase-1), Rsk-2 and Rsk-3, which are phosphorylated by ERK1 and ERK2. Similarly, p38 phosphorylates and activates the serine/threonine kinases MAPKAP kinase-2 and MAPKAP kinase-3 (also designated 3pK). The serine/threonine kinases Mnk1 and Mnk2 are substrates for both ERK and p38 MAP kinases.

REFERENCES

1. Sturgill, T.W., et al. 1988. Insulin-stimulated MAP2 kinase phosphorylates and activates ribosomal protein S6 kinase II. *Nature* 334: 715-718.
2. Stokoe, D., et al. 1992. MAPKAP kinase-2: a novel protein kinase activated by mitogen-activated protein kinase. *EMBO J.* 11: 3985-3994.
3. Davis, R.J. 1993. The mitogen-activated protein kinase signal transduction pathway. *J. Biol. Chem.* 268: 14553-14556.
4. Zhao, Y., et al. 1995. RSK3 encodes a novel pp90rsk isoform with a unique N-terminal sequence: growth factor stimulated kinase function and nuclear translocation. *Mol. Cell. Biol.* 15: 4353-4363.
5. McLaughlin, M.M., et al. 1996. Identification of mitogen-activated protein (MAP) kinase-activated protein kinase-3, a novel substrate of CSBP p38 MAP kinase. *J. Biol. Chem.* 271: 8488-8492.
6. Sithanandam, G., et al. 1996. 3pK, a new mitogen-activated protein kinase-activated protein kinase located in the small cell lung cancer tumor suppressor gene region. *Mol. Cell. Biol.* 16: 868-876.

CHROMOSOMAL LOCATION

Genetic locus: MAPKAPK3 (human) mapping to 3p21.2.

SOURCE

3pK (C-4) is a mouse monoclonal antibody raised against amino acids 1-50 mapping at the N-terminus of 3pK of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

3pK (C-4) is available conjugated to agarose (sc-365148 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-365148 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-365148 PE), fluorescein (sc-365148 FITC), Alexa Fluor® 488 (sc-365148 AF488), Alexa Fluor® 546 (sc-365148 AF546), Alexa Fluor® 594 (sc-365148 AF594) or Alexa Fluor® 647 (sc-365148 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-365148 AF680) or Alexa Fluor® 790 (sc-365148 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

3pK (C-4) is recommended for detection of 3pK of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for 3pK siRNA (h): sc-39105, 3pK shRNA Plasmid (h): sc-39105-SH and 3pK shRNA (h) Lentiviral Particles: sc-39105-V.

Molecular Weight (predicted) of 3pK: 43 kDa.


Molecular Weight (observed) of 3pK: 41 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, HEL 92.1.7 cell lysate: sc-2270 or Jurkat whole cell lysate: sc-2204.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



3pK (C-4): sc-365148. Western blot analysis of 3pK expression in HeLa (A), Jurkat (B), HL-60 (C) and BJAB (D) whole cell lysates. Detection reagent used: m-IgGκ BP-HRP: sc-516102.



3pK (C-4): sc-365148. Near-infrared western blot analysis of 3pK expression in HEL 92.1.7 (A) and HL-60 (B) whole cell lysates. Blocked with UltraCruz® Blocking Reagent: sc-516214. Detection reagent used: m-IgGκ BP-CFL 680: sc-516180.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.