## SANTA CRUZ BIOTECHNOLOGY, INC.

# PYK2 (D-1): sc-365201



#### BACKGROUND

Focal adhesion kinase (FAK) was initially identified as a substrate for the intrinsic protein tyrosine kinase activity of Src-encoded pp60. The deduced amino acid sequence of FAK p125 has shown it to be a cytoplasmic protein tyrosine kinase whose sequence and structural organization are unique compared to other protein families described. A putative new member of the FAK family, designated PYK2 (proline-rich tyrosine kinase 2), exhibits 61% sequence identity with FAK over its kinase domain. PYK2 (also designated CAK $\beta$  or RAFTK) is highly expressed in the central nervous system. Activation of the kinase leads to modulation of ion channel function and the activation of the MAPK signaling pathway. PYK2 is rapidly phosphorylated on tyrosine residues in response to stimuli that increase intracellular calcium levels and compounds that activate members of the PKC family of kinases, such as phorbol esters.

## REFERENCES

- Lipfert, L., et al. 1992. Integrin-dependent phosphorylation of the protein tyrosine kinase pp125<sup>FAK</sup> in platelets. J. Cell Biol. 119: 905-912.
- Guan, J.L. and Shalloway, D. 1992. Regulation of focal adhesion-associated protein tyrosine kinase by both cellular adhesion and oncogenic transformation. Nature 359: 690-692.
- Schaller, M.D., et al. 1992. pp125<sup>FAK</sup>, a structurally distinctive proteintyrosine kinase associated with focal adhesions. Proc. Natl. Acad. Sci. USA 89: 5192-5196.
- Hanks, S.K., et al. 1992. Focal adhesion protein-tyrosine kinase phosphorylated in response to cell attachment to Fibronectin. Proc. Natl. Acad. Sci. USA 89: 8487-8491.

## **CHROMOSOMAL LOCATION**

Genetic locus: PTK2B (human) mapping to 8p21.2; Ptk2b (mouse) mapping to 14 D1.

#### SOURCE

PYK2 (D-1) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 2-27 at the N-terminus of PYK2 of human origin.

## PRODUCT

Each vial contains 200  $\mu g$  lgA kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365201 P, (100  $\mu$ g peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

#### **STORAGE**

Store at 4° C, \*\*D0 NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

# **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### **APPLICATIONS**

PYK2 (D-1) is recommended for detection of PYK2 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for PYK2 siRNA (h): sc-36332, PYK2 siRNA (m): sc-36333, PYK2 shRNA Plasmid (h): sc-36332-SH, PYK2 shRNA Plasmid (m): sc-36333-SH, PYK2 shRNA (h) Lentiviral Particles: sc-36332-V and PYK2 shRNA (m) Lentiviral Particles: sc-36333-V.

Molecular Weight of PYK2: 120 kDa.

Positive Controls: NAMALWA cell lysate: sc-2234, GA-10 whole cell lysate: sc-364230 or Raji whole cell lysate: sc-364236.

## **RECOMMENDED SUPPORT REAGENTS**

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG K BP-HRP: sc-516102 or m-IgG K BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein L-Agarose: sc-2336 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG K BP-FITC: sc-516140 or m-IgG K BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

#### DATA





PYK2 (D-1): sc-365201. Western blot analysis of PYK2 expression in Raji (A), GA-10 (B) and NAMALWA (C) whole cell lysates.

PYK2 (D-1): sc-365201. Immunofluorescence stain ing of methanol-fixed HeLa cells showing nuclear localization.

#### **SELECT PRODUCT CITATIONS**

 Hashimoto-Tane, A., et al. 2016. Micro-adhesion rings surrounding TCR microclusters are essential for T cell activation. J. Exp. Med. 213: 1609-1625.



See **PYK2 (E-3): sc-393181** for PYK2 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor<sup>®</sup> 488, 546, 594, 647, 680 and 790.