

Gas7 (D-2): sc-365372

BACKGROUND

Growth arrest specific proteins, including Gas 1, Gas 6 and Gas 7, are activated in quiescent cells. Gas 7 plays a role in neurite differentiation in cultured mouse cerebellar neurons and PC-12 cells, which makes it a potential therapeutic target to promote the re-establishment of neuronal connections in the injured or disease brain. The gene encoding human Gas7 maps to chromosome 17p13.1, which can translocate with MLL to form MLL-GAS7 fusion products. The Gas7 protein is expressed as three isoforms, a, b, and c, which are differentially expressed in all brain subregions.

REFERENCES

1. Ju, Y.T., et al. 1998. Gas7: a gene expressed preferentially in growth-arrested fibroblasts and terminally differentiated Purkinje neurons affects neurite formation. *Proc. Natl. Acad. Sci. USA* 95: 11423-11428.
2. Lazakovitch, E.M., et al. 1999. The Gas7 gene encodes two protein isoforms differentially expressed within the brain. *Genomics* 61: 298-306.
3. Megonigal, M.D., et al. 2000. Detection of leukemia-associated MLL-GAS7 translocation early during chemotherapy with DNA topoisomerase II inhibitors. *Proc. Natl. Acad. Sci. USA* 97: 2814-2819.
4. She, B.R., et al. 2002. Association of the growth-arrest-specific protein Gas7 with F-Actin induces reorganization of microfilaments and promotes membrane outgrowth. *Exp. Cell Res.* 273: 34-44.
5. Chao, C.C., et al. 2003. Involvement of Gas7 in nerve growth factor-independent and dependent cell processes in PC12 cells. *J. Neurosci. Res.* 74: 248-254.
6. So, C.W., et al. 2003. MLL-Gas7 transforms multipotent hematopoietic progenitors and induces mixed lineage leukemias in mice. *Cancer Cell* 3: 161-171.
7. Lortie, K., et al. 2005. The Gas7 protein potentiates NGF-mediated differentiation of PC12 cells. *Brain Res.* 1036: 27-34.

CHROMOSOMAL LOCATION

Genetic locus: GAS7 (human) mapping to 17p13.1; Gas7 (mouse) mapping to 11 B3.

SOURCE

Gas7 (D-2) is a mouse monoclonal antibody specific for an epitope mapping between amino acids 71-97 near the N-terminus of Gas7 isoform b of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-365372 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

APPLICATIONS

Gas7 (D-2) is recommended for detection of Gas7 of mouse, rat and human origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

Suitable for use as control antibody for Gas7 siRNA (h): sc-45345, Gas7 siRNA (m): sc-45346, Gas7 shRNA Plasmid (h): sc-45345-SH, Gas7 shRNA Plasmid (m): sc-45346-SH, Gas7 shRNA (h) Lentiviral Particles: sc-45345-V and Gas7 shRNA (m) Lentiviral Particles: sc-45346-V.

Molecular Weight of Gas7 isoform a: 38 kDa.


Molecular Weight of Gas7 isoform b: 48 kDa.

Positive Controls: mouse brain extract: sc-2253, mouse cerebellum extract: sc-2403 or PC-12 cell lysate: sc-2250.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Gas7 (D-2): sc-365372. Western blot analysis of Gas7 expression in PC-12 whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.